

EXPRESS MAIL CERTIFICATE

Date: November 23, 1999

Label No. EG 430 341 018 US

I hereby certify that, on the date indicated above I deposited this paper or fee with the U.S. Postal Service and that it was addressed for delivery to the Assistant Commissioner for Patents, Washington, DC 20231 by "Express Mail Post Office to Addressee" service.

A. David Joran  
Registration No. 37,858

Signature

JC675 U.S. PTO  
09/448420

11/22/99

**KRAMER LEVIN NAFTALIS & FRANKEL LLP**

919 Third Avenue  
New York, New York 10022  
212-715-9100

File No: 000973/0004

November 23, 1999

Assistant Commissioner for Patents  
Box Patent Application  
Washington, DC 20231

Sir:

**TRANSMITTAL OF PATENT APPLICATION**

Enclosed please find a Continuation application for United States Letters Patent, based on copending prior PCT International Application No. PCT/US98/10719 which designated the United States, having an International Filing Date of 22 May 1998, claiming priority of U.S. Provisional Application Serial No. 60/047,472, filed May 23, 1997, pursuant to 35 U.S.C. §120 and 37 C.F.R. §1.53(b), as identified below:

Inventors: Michael SEUL, Fanwood, New Jersey; and Richard H. EBRIGHT, North Brunswick, New Jersey.

Title: COLOR-ENCODING AND IN-SITU INTERROGATION OF MATRIX-COUPLED CHEMICAL COMPOUNDS

including the items indicated:

1. Specification and 76 claims: 4 independent; 72 dependent; 1 multiple dependent;

Applicants: Michael SEUL and Richard H. EBRIGHT  
Filed: Concurrently Herewith  
Serial No.: Not Yet Assigned

Atty. Dkt. No.: 000973/0003

2. ☒ Drawings, 9 sheets (Figs. 1-9);
3. ☐ Declaration and power of attorney;
4. ☐ Assignment for recording to: \_\_\_\_\_;
5. ☐ Copy of Statement Claiming Small Entity Status filed in prior application U.S. Serial No. 60/047,472, filed May 23, 1997;
6. ☐ Check in the amount of \$00, (\$ filing; \$ recording);  
☒ See attached Fee Computation Sheet; and
7. ☒ Preliminary Amendment.

Priority is claimed for this application, corresponding application(s) having been filed as follows:


Country: WO  
Number: PCT International Application No. PCT/US98/10719  
International Filing Date: 22 May 1998

Country: US  
Number: 60/047,472  
Date: May 23, 1997

Please amend the specification by inserting before the first line:

--This application is a Continuation of PCT International Application No. PCT/US98/10719, filed May 22, 1998, now pending, which claimed priority of U.S. Provisional Application Serial No. 60/047,472, filed May 23, 1997, now abandoned.--

Respectfully submitted,



A. David Joran  
Registration No. 37,858  
Attorney for Applicants

EXPRESS MAIL CERTIFICATE

Date: November 23, 1999

Label No. EG 430 341 018 US

I hereby certify that, on the date indicated above I deposited this paper or fee with the U.S. Postal Service and that it was addressed for delivery to the Assistant Commissioner for Patents, Washington, DC 20231 by "Express Mail Post Office to Addressee" service.

A. David Joran  
Registration No. 37,858

Signature

Atty. Dkt. No. 000973/0003

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

In re Application of

Michael SEUL and Richard H. EBRIGHT

Serial No: Not Yet Known

Filed: Concurrently Herewith

For: COLOR-ENCODING AND IN-SITU INTERROGATION OF MATRIX-COUPLED  
CHEMICAL COMPOUNDS

Priority International  
Application: PCT/US98/10917

International  
Filing Date: 22 May 1998

-----  
November 22, 1999

Assistant Commissioner for Patents  
Box Patent Application  
Washington, D.C. 20231

Sir:

**PRELIMINARY AMENDMENT**

This Preliminary Amendment is submitted in connection with the filing of a Continuation application identified above based on copending prior PCT International Application No. PCT/US98/10719 which designated the United States.

Prior to examination on the merits, please amend the subject application as follows:

**In the Claims:**

Please amend claims 1, 36, 37, 63-68, and 72 under the provisions of 37 C.F.R. §1.121(c) as follows:

--1. (AMENDED) A method of identifying a compound having a selected property of interest in a library of compounds, each of said compounds being bound to its respective solid support, and being produced by a unique reaction series composed of  $N$  reaction steps, wherein each compound is prepared from a component, and  $N$  is an integer from at least 1 to about 100, which comprises:

- a) dividing a population of solid supports having at least one type of a [first] functional group at the surface of said solid support selected from the group consisting of  $\text{CO}_2\text{H}$ ,  $\text{OH}$ ,  $\text{SH}$ ,  $\text{NH}_2$ ,  $\text{NHR}$ ,  $\text{CH}_2\text{Cl}$ ,  $\text{CH}_2\text{Br}$  and  $\text{CHN}_2$ , wherein  $R$  is a linear  $\text{C}_1\text{-C}_9$  alkyl group, into  $M$  batches, wherein  $M$  is an integer from at least 2 to about 25;
- b) coupling the  $M$  batches of solid support in a set of at least one reaction respectively with  $M$  different components so as to form a bond with the solid support via said [first] functional group, said components being independently protected or unprotected;
- c) adding to each batch, either prior to coupling step b), concurrently therewith, or subsequently to step b), from about 0.001 to about 0.5 molar equivalent of a spectrally distinguishable fluorophore tag associated uniquely with each component, said tag being identified by its characteristic excitation wavelength(s), emission wavelength(s), excited state lifetime and emission intensity, said tag being [activated so as to be] capable of forming [either] a [direct] bond to [the surface of] the solid support[, either via the first or a second functional group which is protected or unprotected and is the same as] or [different from the first functional group bonded] to the component[, or an indirect bond via a  $\text{C}_1\text{-C}_9$  linear or branched alkyl linker moiety which is either interrupted or uninterrupted by at least one oxygen or nitrogen atom or a carbonyl,  $(\text{C}=\text{O})\text{NH}$  or  $\text{NH}(\text{C}=\text{O})$  moiety, wherein when said second functional group is protected, said functional group is deprotected prior to forming said direct or indirect bond, said linker

being bonded to the second functional group at the surface of the solid support]; and  
either

- d) recombining all  $M$  batches, said recombining step being either prior to or subsequent to step e), and steps e) - g); or
- e) performing an assay capable of indicating that any compound in the library either while bound to or cleaved from its solid support has the property of interest;
- f) collecting spectral fluorescence data for each respective solid support so as to determine respective relative abundances of the fluorophore tags bound thereto; and
- g) analyzing the collected spectral fluorescence data by comparing the respective relative abundances of the fluorophore tags determined in step f) so as to determine the unique reaction series for the compound, thereby identifying the compound having the property of interest.--

--36. (AMENDED) An apparatus for identifying a compound having a selected property of interest in a library of compounds, each of said compounds being bound to its respective solid support, and being produced by a unique reaction series composed of  $N$  reaction steps, wherein each said compound is prepared from a component, and  $N$  is an integer from at least 1 to about 100, said solid support being at least one particle array, said apparatus comprising:

- a) an electrode and an electrolyte solution having an interface therebetween,
- b) an electric field generator which generates an electric field at an interface between an electrode and an electrolyte solution;
- c) said electrode being patterned to modify the electrochemical properties of said electrode;
- d) an illuminating source which illuminates said interface with a predetermined light pattern to control the movement of said particles in accordance with said predetermined light pattern and the electrochemical properties of said electrode;
- e) means for preparing said chemical library, which comprises:

- i) means for dividing a population of solid supports having at least one type of a [first] functional group at the surface of said solid support selected from the group consisting of  $\text{CO}_2\text{H}$ ,  $\text{OH}$ ,  $\text{SH}$ ,  $\text{NH}_2$ ,  $\text{NHR}$ ,  $\text{CH}_2\text{Cl}$ ,  $\text{CH}_2\text{Br}$  and  $\text{CHN}_2$ , wherein  $\text{R}$  is a linear  $\text{C}_1$ - $\text{C}_9$  alkyl group, into  $M$  batches, wherein  $M$  is an integer from at least 2 to about 25;
- ii) means for coupling the  $M$  batches of solid support in a set of at least one reaction respectively with  $M$  different components so as to form a bond with the solid support via said [first] functional group, said components being independently protected or unprotected;
- iii) means for adding to each batch either prior to coupling step ii), concurrently therewith, or subsequently to step ii), from about 0.001 to about 0.5 molar equivalent of a spectrally distinguishable fluorophore tag associated uniquely with each component, said tag being identified by its characteristic excitation wavelength(s), emission wavelength(s), excited state lifetime and emission intensity, said tag being [activated so as to be] capable of forming either a [direct] bond to [the surface of] the solid support[, either via the first or a second functional group which is protected or unprotected and is the same as or different from said first functional group, a direct bond] or to the component which if protected is priorly deprotected[, or an indirect bond via a  $\text{C}_1$ - $\text{C}_9$ , linear or branched alkyl linker moiety which is either interrupted or uninterrupted by at least one oxygen or nitrogen atom or a carbonyl,  $(\text{C}=\text{O})\text{NH}$  or  $\text{NH}(\text{C}=\text{O})$  moiety, wherein when said second functional group is protected, said second functional group is deprotected prior to forming said direct or indirect bond, said linker being bonded to said second functional group at the surface of the solid support]; and either
- iv) means for recombining all  $M$  batches, said recombining step either being prior to or subsequent to step v), and steps v)-vii); or;

- v) means for performing an assay capable of indicating that any compound in the library either while bound to or cleaved from its solid support has the property of interest;
- vi) means for collecting spectral fluorescence data for each respective solid support so as to determine respective relative abundances of the fluorophore tags bound thereto;
- vii) means for analyzing the collected spectral fluorescence data by comparing the respective relative abundances of the fluorophore tags determined in step vi) so as to determine the unique reaction series for the compound, thereby identifying the compound having the property of interest.--

--37. (AMENDED) A method of identifying a compound having a selected property of interest in a library of compounds, each of said compounds being bound to its respective solid support, and being produced by a unique reaction series composed of  $N$  coupling or reaction steps, wherein each compound is prepared from components which are independently the same or different, and  $N$  is an integer from at least 1 to about 100, which comprises:

- a) dividing a population of solid supports having at least one type of a [first] functional group at the surface of said solid support surface selected from the group consisting of  $\text{CO}_2\text{H}$ ,  $\text{OH}$ ,  $\text{SH}$ ,  $\text{NH}_2$ ,  $\text{NHR}$ ,  $\text{CH}_2\text{Cl}$ ,  $\text{CH}_2\text{Br}$  and  $\text{CHN}_2$ , wherein  $R$  is a linear  $\text{C}_1\text{-C}_9$  alkyl group, into  $M$  batches, wherein  $M$  is an integer from at least 2 to about 50;
- b) coupling the  $M$  batches of solid support in a set of at least one reaction respectively with  $M$  different initial components so as to form a bond with the solid support via said [first] functional group, said components being protected or unprotected at a group which is capable of participating in a further coupling step and orthogonally protected at non-participating group(s);
- c) adding to each batch either prior to coupling step b), concurrently therewith, or subsequently to step b), from about 0.001 to about 0.5 molar equivalent of a spectrally

distinguishable fluorophore tag associated uniquely with each initial component or a reaction of step b), said tag being identified by its characteristic excitation wavelength(s), emission wavelength(s), excited state lifetime and emission intensity, said tag being [activated so as to be] capable of forming [either] a [direct] bond to [the surface of] the solid support[, either via the first or a second functional group which is protected or unprotected and is the same as or different from said first functional group, a direct bond] or to the initial component [which if protected is priorly deprotected, or an indirect bond via a C<sub>1</sub>-C<sub>9</sub> linear or branched alkyl linker moiety which is interrupted or uninterrupted by either at least one oxygen or nitrogen atom or a carbonyl, (C=O)NH or NH(C=O) moiety, said linker being bonded to said first functional group at the surface of the solid support, wherein when said second functional group is protected, said second functional group is deprotected prior to forming said direct or indirect bond]; and either

d) recombining all  $M$  batches and cleaving any protecting group present at a group which is to participate in a further coupling step, said recombining step being either prior to or subsequent to step e), and steps e)-h); or

e) iteratively  $N - 1$  times

- (1) dividing a population of solid supports into  $M(N)$  batches, wherein  $M(N)$  depends on  $N$  and is an integer from at least 2 to about 25;
- (2) coupling the  $M(N)$  batches of solid support respectively with  $M(N)$  different components, wherein  $M(N)$  is the number of batches during the  $N$ th step, said components being protected or not protected at a group which is capable of participating in a further coupling step and orthogonally protected at a nonparticipating group(s);
- (3) adding to each batch either prior to coupling step (2), concurrently therewith, or subsequently to step (2), from about 0.001 to about 0.5 molar equivalent of a spectrally distinguishable fluorophore tag associated uniquely with each component in the  $N$ th coupling step (2), said tag being identified by its characteristic excitation wavelength(s), emission wavelength(s), excited state



lifetime and emission intensity, said tag being [activated so as to form either] capable of forming a [direct] bond to [the surface of] the solid support[, either via a functional group which is protected or not protected and is the same as or different from the functional group bonded to the component, a direct bond] or to the  $(N - 1)$ th component[, or an indirect bond via a  $C_1 - C_9$  linear or branched alkyl linker moiety which is optionally interrupted by at least one oxygen or nitrogen atom or a carbonyl,  $(C=O)NH$  or  $NH(C=O)$  moiety, said linker being bonded to the functional group at the surface of the solid support, wherein when said functional group is protected, said function group is deprotected prior to forming said direct or indirect bond]; and

(4) recombining all  $M(N)$  batches and cleaving any protecting group present at a group which is to participate in a further coupling step;

so as to form a compound having  $N$  components;

- f) performing an assay capable of indicating that any compound in the library either while bound to or cleaved from its solid support has the property of interest;
- g) collecting spectral fluorescence data for each respective solid support so as to determine respective relative abundances of the fluorophore tags bound thereto; and
- h) analyzing the collected spectral fluorescence data by comparing the respective relative abundances of the fluorophore tags determined in step g) so as to determine the  $N$  components coupled in the unique reaction series for the compound, thereby identifying the compound having the property of interest.--

--63. (AMENDED) The method of claim 62 wherein the dynamic planar array of beads is formed [adjacent to the planar walls of a sandwich flow cell and controlled by light-controlled electrokinetic means] by using an apparatus capable of dynamically assembling an array of beads at an interface between an electrode and an electrolyte solution, said apparatus comprising:

- i) an electrode, an electrolyte solution and an interface therebetween;

- ii) a plurality of beads located in said electrolyte solution;
- iii) said electrode being patterned to include at least one area of modified electrochemical properties; and
- iv) an electric field generator which generates an electric field at said interface to cause the assembly of an array of beads in accordance with the electrochemical properties of said electrode.--

--64. (AMENDED) The method of claim 62 wherein the dynamic planar array of beads is formed by using an apparatus capable of dynamically assembling [and disassembling] an array of beads at an interface between an electrode and an electrolyte solution, said apparatus comprising:

- i) an electrode, an electrolyte solution and an interface therebetween;
- ii) a plurality of beads located in said electrolyte solution;
- iii) [said electrode being patterned to include at least one area of modified electrochemical properties;
- iv)] an illumination source which illuminates said electrode with a predetermined light pattern; and
- [v)] iv) an electric field generator which generates an electric field at said interface to cause the assembly of an array of beads in accordance with the predetermined light pattern.--

--65. (AMENDED) The method of claim 62 wherein spectral fluorescence data are collected for the bead array by initially forming a spatially encoded array of beads suspended at an interface between an electrode and an electrolyte solution, comprising the following steps:

- i) providing an electrode and an electrolyte solution;
- ii) providing multiple types of particles, each type [being stored in accordance with] having chemically or physically distinguishable [particle] characteristics and placing said particles in one [of a plurality of] or more

reservoirs, each reservoir containing [a plurality of like-type] one or more types of said particles suspended in said electrolyte solution;

iii) [providing said reservoirs in the form of an *mxn* grid arrangement] modifying said electrode to define one or more compartments corresponding to one or more of said reservoirs;

iv) [patterning said electrode to define *mxn* compartments corresponding to said *mxn* grid of reservoirs;

v)] depositing [*mxn* droplets] one aliquot from [said *mxn*] one or more of said one or more reservoirs onto said [corresponding *mxn* compartments] modified electrode surface, each said [droplet] aliquot uniquely originating from one of said reservoirs and remaining confined to one of said [*mxn*] one or more compartments and each said [droplet] aliquot containing at least one particle;

[vi)] v) positioning a top electrode above said aliquots so as to simultaneously contact each said [droplet] aliquot;

[vii)] vi) generating an electric field between said top electrode and said [*mxn* droplets] one or more aliquots; and

[viii)] vii) using said electric field to form a bead array in each of said [*mxn*] one or more compartments, each said bead array remaining spatially confined to one of said [*mxn* droplets] one or more aliquots;

ix) illuminating said *mxn* compartments on said patterned electrode with a predetermined light pattern to maintain the position of said bead arrays in accordance with said predetermined light pattern and the pattern of *mxn* compartments; and

x) positioning said top electrode closer to said electrode thereby fusing said *mxn* droplets into a continuous liquid phase, while maintaining each of said *mxn* bead arrays in one of the corresponding *mxn* compartments].--

--66. (AMENDED) The method of claim [65] 62 wherein [said compartments are hydrophilic and the remainder of said electrode surface is hydrophobic] spectral

fluorescence data are collected for the bead array by initially forming a spatially encoded array of beads suspended at an interface between an electrode and an electrolyte solution, comprising the following steps:

- i) providing an electrode and an electrolyte solution;
- ii) providing multiple types of particles, each type having chemically or physically distinguishable characteristics and placing said particles in one or more reservoirs, each reservoir containing one or more types of said particles suspended in said electrolyte solution;
- iii) modifying said electrode to define one or more compartments corresponding to one or more said reservoirs;
- iv) depositing one aliquot from said one or more reservoirs onto said modified electrode surface, each said aliquot uniquely originating from one of said reservoirs and remaining confined to one of said one or more compartments and each said aliquot containing at least one particle;
- v) positioning a top electrode above said aliquots so as to simultaneously contact each said aliquot;
- vi) generating an electric field between said top electrode and said one or more aliquots;
- vii) using said electric field to form a bead array in each of said one or more compartments, each said bead array remaining spatially confined to one of said one or more aliquots;
- viii) positioning said top electrode closer to said electrode thereby fusing said one or more aliquots into a continuous liquid phase, while maintaining each of said one or more bead arrays in one of the corresponding one or more compartments; and
- ix) illuminating said one or more compartments on said patterned electrode with a predetermined sequence of one or more light pattern to place said particle arrays into positions on said electrode surface in accordance with said predetermined sequence of light patterns.--

--67. (AMENDED) [The method of claim 37 wherein  $N$  is an integer from at least 3 to about 12] A planar array encoded in accord with claim 62.--

--68. (AMENDED) [The method of claim 37 wherein  $M$  and  $M(N)$  are independently an integer from at least 4 to about 12] A planar array encoded in accord with claim 63, 64, 65 or 66.--

--72. (AMENDED) An apparatus for identifying a compound having a selected property of interest in a library of compounds, each of said compounds being bound to its respective solid support, and being produced by a unique reaction series composed of  $N$  coupling and reaction steps, wherein each said compound is prepared from a set of components which are independently the same or different, and  $N$  is an integer from at least 1 to about 100, said solid support being at least one particle array, said apparatus comprising:

- a) an electrode and an electrolyte solution having an interface therebetween;
- b) an electric field generator which generates an electric field at an interface between an electrode and an electrolyte solution;
- c) said electrode being patterned to modify the electrochemical properties of said electrode;
- d) an illuminating source which illuminates said interface with a predetermined light pattern to control the movement of said particles in accordance with said predetermined light pattern and the electrochemical properties of said electrode;
- e) means for preparing said chemical library, which comprises:
  - i) means for dividing a population of solid supports having at least one type of a [first] functional group at the surface of said solid support selected from the group consisting of  $\text{CO}_2\text{H}$ ,  $\text{OH}$ ,  $\text{SH}$ ,  $\text{NH}_2$ ,

NHR, CH<sub>2</sub>Cl, CH<sub>2</sub>Br and CHN<sub>2</sub>, wherein R is a linear C<sub>1</sub>-C<sub>9</sub> alkyl group, into *M* batches, wherein *M* is an integer from at least 2 to about 50;

ii) means for coupling the *M* batches of solid support in a set of at least one reaction respectively with *M* different initial components so as to form a bond with the solid support via said [first] functional group, said components being protected or unprotected at a group which is to participate in a further coupling step and orthogonally protected at non-participating group(s);

iii) means for adding to each batch either prior to coupling step ii), concurrently therewith, or subsequently to step ii), from about 0.001 to about 0.5 molar equivalent of a spectrally distinguishable fluorophore tag associated uniquely with each initial component, said tag being identified by its characteristic excitation wavelength(s), emission wavelength(s), excited state lifetime and emission intensity, said tag being [activated so as to be] capable of forming [either] a [direct] bond to [the surface of] the solid support[, either via the first or a second functional group which is protected or unprotected and is the same as] or [different from said first functional group bonded] to the component[, or an indirect bond via a C<sub>1</sub>-C<sub>9</sub>, linear or branched alkyl linker moiety which is either interrupted or uninterrupted by either at least one oxygen or nitrogen atom or a carbonyl, (C=O)NH or NH(C=O) moiety, said linker being bonded to said second functional group at the surface of the solid support, wherein when said second functional group is protected, said second functional group is deprotected prior to forming said direct or indirect bond]; and either

iv) means for recombining all *M* batches and cleaving any protecting group present at a group which is to participate in a further coupling step, and steps v)-viii); or

v) means for iteratively *N* - 1 times

(1) dividing a population of solid supports into  $M(N)$  batches, wherein  $M(N)$  depends on  $N$  and is an integer from at least 2 to about 25;

(2) coupling the  $M(N)$  batches of solid supports respectively with  $M(N)$  different components, wherein  $M(N)$  is the number of batches during the  $N$ th step, said components being protected or unprotected at a group which is capable of participating in a further coupling step and orthogonally protected at a non-participating group(s);

(3) adding to each batch either prior to coupling step (2), concurrently therewith, or subsequently to step (2), from about 0.001 to about 0.1 molar equivalent of a different spectrally distinguishable fluorophore tag associated uniquely with each component during the  $N$ th coupling step (2), said tag being uniquely identified by its excitation wavelength, emission wavelength, excited-state lifetime or emission intensity, whereby said tag is [activated so as to be] capable of forming [either] a [direct] bond to the solid support[, either via an  $N$ th functional group which is protected or unprotected and is the same as or different from the first functional group, or an indirect bond thereto via a  $C_1$ - $C_9$  linear or branched alkyl linker moiety which is either interrupted or uninterrupted by either at least one oxygen or nitrogen atom or a carbonyl or  $NH(C=O)$  moiety,] or [a direct bond] to the  $(N-1)$ th component which if protected is priorly deprotected[, said tag or linker being bound via the group which is to participate in a further coupling step, wherein when said  $N$ th functional group is protected, said  $N$ th functional group is deprotected prior to forming said direct or indirect bond]; and

(4) recombining all  $M(N)$  batches and

cleaving the protecting group present if present at a group which is to participate in a further coupling step;

so as to form a compound having  $N$  components;

vi) means for performing an assay capable of indicating that any compound in the library either while bound to or cleaved from its solid support has the property of interest;

vii) means for collecting spectral fluorescence data for each respective solid support so as to determine respective relative abundances of the fluorophore tags bound thereto;

viii) means for analyzing the collected spectral fluorescence data by comparing the respective relative abundances of the fluorophore tags determined in step vii) so as to determine the  $N$  components coupled in the unique reaction series for the compound, thereby identifying the compound having the selected property of interest.-

Please add new claim 73 as follows:

--73. (NEW) An apparatus for identifying a compound having a selected property of interest in a library of compounds, each of said compounds being bound to its respective solid support, and being produced by a unique reaction series composed of  $N$  coupling and reaction steps, wherein each said compound is prepared from a set of components which are independently the same or different, and  $N$  is an integer from at least 1 to about 100, said solid support being at least one particle array, said apparatus comprising at least one particle array prepared in accordance with claim 62, and further comprising means for preparing said chemical library, which comprises:

i) means for dividing a population of solid supports having at least one type of a functional group at the surface of said solid



support selected from the group consisting of  $\text{CO}_2\text{H}$ ,  $\text{OH}$ ,  $\text{SH}$ ,  $\text{NH}_2$ ,  $\text{NHR}$ ,  $\text{CH}_2\text{Cl}$ ,  $\text{CH}_2\text{Br}$  and  $\text{CHN}_2$ , wherein  $\text{R}$  is a linear  $\text{C}_1\text{-C}_9$  alkyl group, into  $M$  batches, wherein  $M$  is an integer from at least 2 to about 50;

ii) means for coupling the  $M$  batches of solid support in a set of at least one reaction respectively with  $M$  different initial components so as to form a bond with the solid support via said functional group, said components being protected or unprotected at a group which is to participate in a further coupling step and orthogonally protected at non-participating group(s);

iii) means for adding to each batch either prior to coupling step ii), concurrently therewith, or subsequently to step ii), from about 0.001 to about 0.5 molar equivalent of a spectrally distinguishable fluorophore tag associated uniquely with each initial component, said tag being identified by its characteristic excitation wavelength(s), emission wavelength(s), excited state lifetime and emission intensity, said tag being capable of forming a bond to the solid support or to the component; and either

iv) means for recombining all  $M$  batches and cleaving any protecting group present at a group which is to participate in a further coupling step, and steps v)-viii); or

v) means for iteratively  $N - 1$  times

(1) dividing a population of solid supports into  $M(N)$  batches, wherein  $M(N)$  depends on  $N$  and is an integer from at least 2 to about 25;

(2) coupling the  $M(N)$  batches of solid supports respectively with  $M(N)$  different components, wherein  $M(N)$  is the number of batches during the  $N$ th step, said components being protected or unprotected at a group which is capable of participating in a further coupling step and orthogonally protected at a non-participating group(s);

(3) adding to each batch either prior to coupling step (2), concurrently therewith, or subsequently to step (2), from about 0.001 to about 0.1 molar equivalent of a different spectrally distinguishable fluorophore tag associated uniquely with each component during the  $N$ th coupling step (2), said tag being uniquely identified by its excitation wavelength, emission wavelength, excited-state lifetime or emission intensity, whereby said tag is capable of forming a bond to the solid support or to the  $(N-1)$ th component which if protected is priorly deprotected; and

(4) recombining all  $M(N)$  batches and cleaving the protecting group present if present at a group which is to participate in a further coupling step;

so as to form a compound having  $N$  components;

vi) means for performing an assay capable of indicating that any compound in the library either while bound to or cleaved from its solid support has the property of interest;

vii) means for collecting spectral fluorescence data for each respective solid support so as to determine respective relative abundances of the fluorophore tags bound thereto;

viii) means for analyzing the collected spectral fluorescence data by comparing the respective relative abundances of the fluorophore tags determined in step vii) so as to determine the  $N$  components coupled in the unique reaction series for the compound, thereby identifying the compound having the selected property of interest.-

In re Application of Michael SEUL and Richard H. EBRIGHT  
Serial No: Not Yet Known  
Filed: Concurrently Herewith

000973/003

**REMARKS**

The subject application is a Continuation Application of copending PCT International Application No. PCT/US98/10719, which claims the benefit of U.S. Provisional Application Serial No. 60/047,472, filed May 23, 1997.


Claims 1-72 were pending in the prior International application. Claims 1, 36, 37, 63-68, and 72 have been amended herein, and new claim 73 is added herein.. Accordingly, claims 1-73 are under examination. It is maintained that the amendments set forth herein merely make more clear what is applicants' intended invention. The amendments do not introduce an issue of new matter, and therefore, applicants respectfully request entry of the claims as amended.

In view of the amendments and remarks made herein, the claims are believed to be in allowable condition, and prompt examination on the merits is requested.

If a telephone interview would be of assistance in advancing prosecution of the subject application, applicants' undersigned attorney invites the Examiner to telephone him at the number provided.

No fee, aside from the initial filing fees and the fee for new claim 73, payment for which will be made in due course, is believed necessary in connection with the filing of this Preliminary Amendment. However, if any fee is otherwise deemed required, authorization is hereby given to charge the amount of such fee to Deposit Account 50-0540.

Respectfully submitted,



A. David Joran  
Registration No. 37,858  
Attorney for Applicants

KRAMER, LEVIN, NAFTALIS & FRANKEL LLP  
919 Third Avenue  
New York, N.Y. 10022  
(212) 715-9100  
(212) 715-8000 (facsimile)

PRELIMINARY  
AMENDMENT

Page 17

## Color-Encoding and In-situ Interrogation of Matrix-Coupled Chemical Compounds

5

### Field of the Invention

The present invention generally relates to the field of analytical chemistry.

The present invention specifically relates to a highly parallel mode of presenting and  
10 probing multiple chemical compounds, with applications to combinatorial library synthesis,  
ultrahigh-throughput screening, diagnostic assays for multiple agents and sensors. The  
present invention introduces several color codes to label collections of carrier particles such  
as colloidal beads; in addition, the present invention describes a method and apparatus for the  
in-situ interrogation of beads or collections of beads by way of multi-color fluorescence  
15 imaging and spectral analysis of individual beads to ascertain the chemical identities of bead-  
anchored compounds. The encoding of beads by simple and extended simple color codes and  
by binary and extended binary color codes may be augmented by measuring bead size and  
shape or other physico-chemical properties such as polarizability embedded in the bead core.

### 20 Background of the Invention

#### *1-Solid Phase Chemical Libraries*

An emerging paradigm for lead discovery in pharmaceutical and related industries  
such as agricultural biotechnology, is the assembly of novel synthetic compound libraries by  
new methods of solid state "combinatorial" synthesis. Combinatorial chemistry refers to a set  
25 of strategies for the parallel synthesis and testing of multiple compounds or compounds  
mixtures, either in solution or in solid supports in the form of beaded resins ("beads"). In  
general, a combinatorial synthesis employing M precursors in each of N reaction steps

produces  $M^N$  compounds. For example, a combinatorial synthesis produces  $4^N$  oligonucleotides in  $N$  steps, each employing 4 oligonucleotide precursors; similarly, a combinatorial synthesis of  $N$  steps, each employing 20 amino acid precursors, produces  $20^N$  oligopeptides.

5

### *1.1 - One Bead/One Compound Chemical Libraries*

One implementation of combinatorial synthesis that is suitable to produce very large chemical libraries relies on solid supports in the form of beaded resins ("beads") and encodes reaction steps in a "divide, couple and recombine" (DCR) strategy (Fig. 1), also referred to as "resin-splitting" synthesis. The resulting "one bead/one compound" chemical libraries contain from  $10^6$  to  $10^8$  compounds. These libraries are screened by performing a wide variety of chemical and biochemical assays to identify individual compounds eliciting a positive response. The chemical identity of such compounds can be determined by direct analysis.

15 Two methods of direct analysis are micro-sequencing and mass spectrometry. Both methods require the physical isolation of synthesis beads displaying compounds of interest and both require off-line chemical analysis based on substantial amounts of compound - tens to hundreds of picomoles. Micro-sequencing, limited to libraries of oligopeptides and oligonucleotides, does not distinguish between stereoisomers. Mass spectrometry is unable to distinguish between precursors of equal mass such as D- and L-amino acids or leucine and isoleucine. The requirement of direct chemical analysis for a substantial quantity of compound dictates the use of large bead resins (a typical bead diameter is  $130\mu\text{m}$ ) to ensure that picomolar quantities of each compound can be recovered, even when it is becoming increasingly desirable to perform high throughput screening of the compound library in miniaturized environments to reduce requisite volumes of sample and reagents and to enhance throughput.

25

### *1.2 - Encoded One Bead/One Component Chemical Libraries*

One approach to overcoming the serious limitations of standard one bead/one compound chemical libraries is to encode chemical compound identities. This facilitates the identification of compounds not amenable to direct determination by micro-sequencing or mass spectrometry. One encoding method employs the co-synthesis of peptides and

30

oligonucleotides to represent the identity of non-sequenceable synthesis products (Nikolaiev et al., "Peptide-Encoding for Structure Determination of Non-Sequenceable Polymers Within Libraries Synthesized and Tested on Solid-Phase Supports", *Peptides Res.* 6, 161 (1993), the contents of which are included herein by reference). A second method, compatible with a wider range of chemical reaction conditions, employs a set of tagging molecules to record the reaction histories of beads.

One implementation of the latter method uses a set of pre-synthesized, chromatographically distinguishable molecular tags T1, T2,..., TM to construct a chemical binary code. In prior art, molecular tags are structurally related molecules (Fig. 2) which can be identified by their characteristic gas chromatographic retention times (Still et al., "Complex combinatorial libraries encoded with tags", U.S. Patent No. 5,565,324, the contents of which are included herein by reference).

At each step of DCR synthesis, a unique tag from the set is added to each divided aliquot to record the reaction carried out with that aliquot. The concept may be illustrated by examining the steps of a 2-step synthesis using reagents  $R^1_1$ ,  $R^1_2$  and  $R^1_3$  in step 1, and reagents  $R^2_1$ ,  $R^2_2$  and  $R^2_3$  in step 2, to generate nine products. The reagents of the first step are uniquely identified by the binary addresses 01 ( $R^1_1$ ), 10 ( $R^1_2$ ) and 11 ( $R^1_3$ ), and the reagents of the second step are uniquely identified by the binary addresses 01 ( $R^2_1$ ), 10 ( $R^2_2$ ) and 11 ( $R^2_3$ ). Each binary address is chemically represented in terms of a set of molecular tags: T1 (01 in step 1 representing  $R^1_1$ ), T2 (10 in step 1 representing  $R^1_2$ ) and T2T1 (11 in step 1 representing  $R^1_3$ ) and analogously with T3 (01 in step 2 representing  $R^2_1$ ), T4 (10 in step 2 representing  $R^2_2$ ) and T4T3 (11 in step 2 representing  $R^2_3$ ).

A sequence of reaction steps is recorded by simply concatenating binary addresses. Thus, 11.01, read right to left, would indicate the sequence "reagent  $R^2_3$  in step 2, reagent  $R^1_1$  in step 1". The chemical representation of this sequence is T4T3.T1, and the presence on the bead of this particular set of tags indicates the chemical identity of the bead-anchored synthesis product. The strategy is readily generalized to larger reactions. For example, 7 reagents to be used in each reaction step can be uniquely identified by the binary addresses 001 ( $R^1_1$ ), 010 ( $R^1_2$ ), ..., 111 ( $R^1_7$ ). Although superior to un-encoded one bead/one compound methods, nevertheless the tagging strategy of prior art still suffer from three limitations. First, individual beads of interest must be physically isolated from the rest; next, molecular tags

must be chemically or photochemically cleaved from the bead and cleaved tags must be collected; and finally, chemical analysis (e.g., gas chromatography) must be performed. These numerous time- and labor-intensive manipulations eliminate much of the enhancement in throughput gained by the DCR synthesis strategy.

5

### *1.3 Screening and Lead Compound Optimization*

The high specificity of typical biological substrate-target interactions implies that the vast majority of compounds in a library will be inactive for any particular target. Thus, the task of screening is to identify the very few compounds within the library that display activity  
10 in binding or in functional assays. Common targets include enzymes and receptors as well as nucleic acids.

To implement the rapid screening and scoring of an entire library of synthetic compounds, in practice containing  $10^4$  to  $10^8$  compounds, requires systematic screening procedures if the task is to be completed within viable time frames. Several assay formats  
15 have been described to implement the screening of bead-based combinatorial libraries. These include: reaction of a collection of beads, allowed to settle under gravity, with an enzyme-labeled or fluorophore-labeled target molecule followed by visual detection (Lam et al., "A new type of synthetic peptide library for identifying ligand-binding activity", *Nature* 354 (1991), the contents of which are included herein by reference); incubation of beads with  
20 radio-labeled target molecules and subsequent agarose immobilization of beads and autoradiographic detection (Kassarjian, Schellenberger and Turck, "Screening of Synthetic Peptide Libraries with Radio-labeled Acceptor Molecules", *Peptide Res.* 6, 129 (1993), the contents of which are included herein by reference); and partial release of compounds from beads for solution-phase testing (Salmon et al., "Discovery of biologically active peptides in random  
25 libraries: Solution-phase testing after staged orthogonal release from resin beads", *Proc. Natl. Acad. Sci. USA* 90, 11708 (1993), the contents of which are included herein by reference).

WO95/32425 provides a method of preparing combinatorial libraries using a method of encoding combinatorial libraries with fluorophore labeled beads. According to the method, a first combinatorial library is prepared by conducting a set of reactions on tagged beads to  
30 afford an encoded first registry (i.e., step in the synthetic sequence). A second combinatorial library is prepared using similar reaction steps but the tagged beads are combined and

separated prior to the first reaction sequence and the beads are sorted prior to the second reaction sequence. Subsequent libraries are prepared as for the second library except that the sorting step takes place prior to a different registry in each subsequent library. Thus, WO95/32425 teaches only individually labelling the first step and physical separation of beads to identify each modified combinational library.

Nederlof et al., *Cytometry*, 13, 839-845 (1992), teaches the use of ratio labeling as a way of increasing the number of simultaneously detectable probes beyond the seven used previously. In this approach, ratio-labelled probes are identified on the basis of the ratio of color intensity, not just the particular colors used. Fluorescence ratios are measured and used as additional encoding colors. The method requires double-labeling of probes using different ratios of labels. The method is not specifically directed to synthetic combinational libraries. Accordingly, the field of Nederlof's method is the detection of multiple DNA/RNA sequence by in situ hybridization, and is not relevant to the field of encoding of synthetic chemical libraries.

Speiche, Ballard & Ward, *Nature Genetics*, 12, 368 (1996), describe a method of characterizing complex chromosomal karyo types using multi-fluorescence in situ hybridization. Instead of using ratio-double labelling as in Nederlof, Speiche et al. use a set of six fluorescent dyes with spectral emission peaks spread across the photometric response range to visualize 27 combinationally labelled probes. Speiche et al. do not disclose a method of encoding synthetic combinational libraries.

Still et al., *Proc. Nat'l Acad. Sci.*, 90, 10922-926 (1993), disclose a method of synthesis of tagged combinational libraries using a binary code based on different electrophoretic tags. The method requires use of photocleavable molecular tags which comprise variously substituted aryl moieties linked via a variable-length aliphatic hydrocarbon chain, whereby the tags when cleaved are distinctly resolvable by capillary gas chromatography with electrochemical detection. Color detection is not used in this method. The method also requires cleavage from the solid support in order to analyze the sequence. In related work, Still et al. U.S. 5,721,099 disclose methods of preparing encoded combinatorial libraries, but again the method requires cleavage of the identifier tags prior to analysis of the encoded reaction history. In contrast, the present invention provides an in situ approach to the interrogation of encoded combinatorial libraries, and represents an advance over the prior



methods of encoding libraries. The success of the present invention is unexpected in view of the prior approaches because of the scattering phenomena expected for a spectral analysis performed in heterogeneous media which would dissipate spectral signal-to-noise giving rise to practical difficulties in detecting accurately relative abundance information for fluorophore tags. The present methodology demonstrates for the first time a way of solving these practical problems in performing in situ encoding and interrogation of combinatorial libraries .

## *II - Multi-Agent Monitoring and Diagnostics*

Diagnostic panels display multiple chemistries to screen unknown solutions for the presence of multiple agents. For example, blood group specificity is determined by spotting an unknown blood sample onto a panel of surface-bound antibodies whose arrangement in the panel reflects their antigen-specificity. Antigen-binding to any specific patch in the panel reveals the chemical identity of the antigen and enhance the blood type. Another realization of the same concept of displaying multiple diagnostic probes in a spatially encoded panel or array involves screening of mutations by assaying for hybridization of DNA to one of a large number of candidate matching strands which are placed in known positions on a planar substrate in a checkerboard pattern. This may be achieved by dispensing droplets containing distinct probes, or may involve the in-situ synthesis of oligonucleotide strands of varying composition.

Spatial encoding relies on the panel or array fabrication process to preserve chemical identity, adding time and expense. As the number of fields in the checkerboard increases, so does the challenge of fabricating the requisite array. In addition, probes must be immobilized - usually by adhesion to the surface of a planar substrate - to maintain the integrity of the spatial encoding scheme. In practice, this assay format can be problematic: sample accumulation can be slow and probe accessibility restricted.

## *III - Current Applications of Multicolor Fluorescence Detection*

The present invention describes a method and apparatus for in-situ interrogation and deconvolution of bead-based combinatorial libraries using multi-color fluorescence imaging and spectral analysis. Recent applications of multi-color fluorescence spectroscopy to DNA sequencing and chromosome painting place requirements on sensitivity and wavelength

selectivity exceeding those encountered in conventional applications such as determinations of fluorescence intensity ratios.

Within the context of DNA sequencing, a variety of configurations for rapid detection of 4-color fluorescence have been described. These involve: a dedicated photomultiplier tube  
5 detector for each emission wavelength, with corresponding sets of beam splitters in the optical path to produce spatially separated beams; a single detector and rotating filterwheel to select the desired set of wavelengths in a multiplexed recording mode; or a dispersive arrangement that relies on a prism or grating to split the emitted light from multiple fluorophores according to wavelength and takes advantage of recent advances in charge-coupled device (CCD)  
10 technology to record spectra on an integrating linear or rectangular CCD array (Karger et al., "Multiwavelength fluorescence detection for DNA sequencing using capillary electrophoresis", Nucl. Acids Res. 19, 4955 (1991), the contents of which are incorporated herein by reference).

## 15 **Summary of the Invention**

The present invention provides a method to construct several color codes for the purpose of uniquely labeling members of a group of beads or equivalent objects ("beads") to preserve the chemical identity of the beads and thus the identity of bead-coupled chemical compounds. These color codes are based on a set of encoding fluorophores of distinguishable  
20 wavelengths, excited-state lifetimes and levels of intensity, the latter controlled by adjusting the abundances of dyes. Specifically, the present invention describes a method and apparatus for the encoding and in-situ interrogation of a set of distinct, bead-based chemistries.

Binary and extended binary color codes offer large coding capacity and represent a general strategy to encode multi-step reaction histories such as those encountered in divide-  
25 couple-recombine (DCR) synthesis strategies for combinatorial chemical libraries, as illustrated and discussed herein.

Simple and extended simple color codes offer an efficient strategy to encode a smaller set of distinct chemistries that are typical of panels displaying multiple targets or probes in biochemical assays including multi-agent diagnostic and environmental tests and other  
30 biochemical assays.

All color codes can be augmented by varying distinguishable features of beads such as shape and size or other suitable physico-chemical parameter associated with bead cores such as polarizability.

The identity of the compound anchored to any specific bead is determined in-situ by optically probing individual beads to read the color code, as described herein. This ensures the identification of bead-anchored chemical compounds without the need for physical separation and without the need for off-line chemical analysis.

The encoding strategy of the present invention is compatible with all formats of bead-based combinatorial synthesis and screening described to date. A preferred implementation that has the advantage of enabling miniaturization and automation of screening and decoding operations relies on planar bead arrays which may be formed, maintained and manipulated adjacent to a planar electrode surface.

### Brief Description of the Drawings

Other objects, features and advantages of the invention discussed in the above brief explanation will be more clearly understood when taken together with the following detailed description of an embodiment which will be understood as being illustrative only, and the accompanying drawings reflecting aspects of that embodiment, in which:

Fig. 1 is an illustration of "Divide-Couple-Recombine" combinatorial synthesis;

Fig. 2 is an illustration of labeling individual synthesis beads with chemical tags ("bar codes"). Examples of molecular structures used for such tags are also shown: different tags are made by varying  $n$  and  $Ar$ ;

Fig. 3 is an illustration of two alternative methods of placing fluorophore or chromophore tags (F) on synthesis beads;

Fig. 4 is an illustration of binary color coding with fluorophores, Y, B, G and R. The example enumerate coded bead populations produced in combinatorial peptide synthesis employing reagents  $R^1_1, R^1_2, R^1_3$  and  $R^1_4$  in step 1 and reagents  $R^2_1, R^2_2, R^2_3$  and  $R^2_4$  in step 2 (see also: Table I);

Fig. 5 is an illustration of emission spectra of the CyDye family of commercially available fluorescent dyes whose spectral characteristics are summarized in the table

accompanying the figure (Amersham LIFE SCIENCE, Catalog of Multicolor Fluorescent Reagents, 1995, the contents of which are included herein by reference);

Fig. 6 is an illustration of a random bead array encoded according to the simple color code  $SCC(l=1, m=5)$ ;

5 Fig. 7 is an illustration of a multi-color fluorescence microscope with integrated spectral analysis based on dispersive optics;

Fig. 8 is an illustration of several geometries of multi-color fluorescence imaging and spectrometry.

10 Fig. 9 is an illustration of an example of a solid support having a hydroxy functional group at its surface which is modified by a linker which is formed in a multistep process involving a deprotection of an Mmt protecting group and subsequent reaction with an activated ester of a fluorescent dye in accord with the present invention.

## Detailed Description of the Preferred Embodiment

### 15 *Implementation of Color Codes*

The color coding strategy of the present invention provides a method to place a set of fluorophores - or, more generally, chromophores - on each bead so as to uniquely encode the chemical identity of the compound on that bead. Specifically, during each coupling step in the course of DCR combinatorial synthesis, one or more fluorophores are attached to each  
20 bead. Decoding is based on the determination of relative abundances of fluorophores on a bead of interest by in-situ optical interrogation.

Fluorophores can be added in two ways. In the first method, the fluorophore is added directly to a small fraction of the nascent compound, thereby terminating further synthesis of that fraction of nascent compound (Fig. 3A). In the second method, the label is covalently  
25 attached to reserved reaction sites other than nascent compound to ensure that precursors are not terminated by labeling (Fig. 3B). In the first method and in most implementations of the second method, the quantity,  $x$ , of fluorophore added to each bead is sub-stoichiometric with respect to nascent compound, with  $x$  typically in the range 0.001 to 0.1 mole equivalents of nascent compound on the bead. Three factors govern the choice of  $x$ . First, the density of  
30 tags on beads must not materially interfere with synthesis and with subsequent screening assays. Second, the density of tags on beads must remain sufficiently low as to avoid

complication due to fluorescence energy transfer. Third, labeled sites must be present in sufficient number to meet the requirements of signal detection and discrimination, as discussed herein.

To implement the color coding strategy, the present invention takes advantage of three properties of fluorophores to construct an alphabet of fluorophore tags, namely: emission wavelength; excited-state lifetime; and emission intensity. Denoting by  $m_F$  the number of available fluorophores with distinguishable emission maxima and/or excited state lifetimes, and denoting by  $m_I$  the number of distinguishable intensity levels, controlled by adjusting relative quantities of fluorophores (e.g. x, 2x, 3x..), the size of the alphabet of fluorophore tags is  $m = m_F \cdot m_I$ . The surfaces of labeled beads will display a multiplicity of distinct fluorophores (see Fig. 4). In-situ optical interrogation of these multi-colored beads serves to record emission spectra from which relative abundances of fluorophores are determined to decipher the color code, as discussed and illustrated herein.

### Binary Color Codes

One rendition of this code is a binary color code (BCC) using  $m_F$  fluorophores, all with  $m_I = 1$ . This BCC will encode up to  $2^{m_F}$  distinct compounds. In this BCC, the  $m_F$  fluorophores could differ in excite-state lifetimes, emission maxima or both. For convenience, the following specific example uses fluorophores differing solely in their emission maxima ("colors"). The combinatorial synthesis of 16 products in two reaction steps, each using a set of  $N=4$  reagents, would be encoded as follows:

**Table I**

25	Step 1: $R^1_1$ (00)	No color	$R^1_2$ (01) Red	$R^1_3$ (10) Green	$R^1_4$ (11) Red+Green
	Step 2: $R^2_1$ (00)	No color	$R^2_2$ (01) Blue	$R^2_3$ (10) Yellow	$R^2_4$ (11) Yellow+Blue
30	$R^2_1, R^1_1$	00.00 NN.NN	no color	$R^2_3, R^1_1$ 10.00	YN.NN Y
	$R^2_1, R^1_2$	00.01 NN.NR	R	$R^2_3, R^1_2$ 10.01	YN.NR YR
	$R^2_1, R^1_3$	00.10 NN.GN	G	$R^2_3, R^1_3$ 10.10	YN.GN YG
	$R^2_1, R^1_4$	00.11 NN.GR	GR	$R^2_3, R^1_4$ 10.11	YN.GR YGR
	$R^2_2, R^1_1$	01.00 NB.NN	B	$R^2_4, R^1_1$ 11.00	YB.NN YB

$R^2_2, R^1_2$	01.01 NB.NR	BR	$R^2_4, R^1_2$	11.01	YB.NR	YBR
$R^2_2, R^1_3$	01.10 NB.GN	BG	$R^2_4, R^1_3$	11.10	YB.GN	YBG
$R^2_2, R^1_4$	01.11 NB.GR	BGR	$R^2_4, R^1_4$	11.11	YB.GR	YBGR

5 The binary representation of four reagents is  $R_1(00)$ ,  $R^1_2(01)$ ,  $R^1_3(10)$  and  $R^1_4(11)$  for the reagents used in step 1, and  $R^2_1(00)$ ,  $R^2_2(01)$ ,  $R^2_3(10)$  and  $R^2_4(11)$  for those in step 2. As before, sequences of reaction steps correspond to concatenated binary codes, and in the example all  $4^2=16$  possible sequences are represented by 4-bit strings. Thus, the sequence: "reagent  $R^2_3$  in step 2, reagent  $R^1_4$  in step 1" would be represented by the string 10.11 (read  
10 right to left). Using an alphabet of four fluorophores, with colors denoted by R, G, B, and Y as before, and assigned (Y, B, G, R) to represent 4-bit strings, the  $2^4$  possible strings (read right to left) are encoded in BCC ( $m=4$ ) as displayed in table I and in Fig. 4.

A second rendition of the color code is a binary color code using  $m_F$  fluorophores with varying relative abundances and thus varying intensities at each step. The resulting eXtended  
15 binary color code (XBCC) will encode  $2^{(m_F \cdot m_I)}$  distinct compounds. For example, using an alphabet (2G, 2R, G, R) with only two distinct colors to represent 4-bit strings,  $2^4$  possible strings (read right to left) are encoded in XBCC ( $m_F=2$ ,  $m_I=2$ ) as enumerated in Table II. In the example, deconvolution will require discrimination of four distinct intensity levels for each of the two emission bands. If  $N$  steps are involved, the number of intensity  
20 levels to be discriminated in the extended binary color code XBCC ( $m_F$ ,  $m_I$ ) may be as high as  $N \cdot m_I$ . The attainable intensity discrimination is ultimately limited by the signal-to-noise ratio attainable in the spectral analysis of individual beads.

25 **Table II**

30	Step 1: $R^1_1(00)$	No color	$R^1_2(01)$ Red	$R^1_3(10)$ Green	$R^1_4(11)$ Red+Green
	Step 2: $R^2_1(00)$	No color	$R^2_2(01)$ 2Red	$R^2_3(10)$ 2Green	$R^2_4(11)$ 2Red+2Green
35	$R^2_1, R^1_1$	00.00 NN.NN	no color	$R^2_3, R^1_1$	10.00 2GN.NN GG
	$R^2_1, R^1_2$	00.01 NN.NR	R	$R^2_3, R^1_2$	10.01 2GN.NR GGR
	$R^2_1, R^1_3$	00.10 NN.GN	G	$R^2_3, R^1_3$	10.10 2GN.GN GGG
	$R^2_1, R^1_4$	00.11 NN.GR	GR	$R^2_3, R^1_4$	10.11 2GN.GR GGGR

$R^2_2, R^1_1$	01.00 N2R.NN	RR	$R^2_4, R^1_1$	11.00	2G2R.NN	GGRR
$R^2_2, R^1_2$	01.01 N2R.NR	RRR	$R^2_4, R^1_2$	11.01	2G2R.NR	GGRRR
$R^2_2, R^1_3$	01.10 N2R.GN	RRG	$R^2_4, R^1_3$	11.10	2G2R.GN	GGGRR
$R^2_2, R^1_4$	01.11 N2R.GR	RRRG	$R^2_4, R^1_4$	11.11	2G2R.GR	GGGRRR

5

Another example describes the color-coding of products created in a combinatorial synthesis using 7 reagents in the first step, 6 reagents in each of the final two steps.

Reagents are represented by binary addresses R1(001), R2(010), R3(011)...R7(111); for simplicity of notation, we omit the superscript for reagents (R) used in different steps.

10 Let  $m_F=4$  (color denoted as before) and  $m_I=2$ . The following XBCC based on an 8-letter alphabet (2Y, 2B, 2G, 2R, Y, B, G, R) and illustrated in Table III may be devised to encode the  $7 \cdot 6 \cdot 6 = 252$  synthesis products created in this synthesis. While the construction of the XBCC would require 9-bit strings to represent the full set of  $8^3 = 512 = 2^9$  configurations created by all possible concatenations of 3-bit strings, the actual 252 required configurations of the example can in fact be accommodated in the set of  $2^8$  possible 8-bit strings by making replacements of the sort indicated in the example. Thus, the reaction sequence "reagent 6 in step 3, reagent 1 in step 2, reagent 3 in step 1" is represented by the XBCC ( $m_F=4, m_I=2$ ) as follows (read right to left): R6.R1.R3 = 2X2B.N.G = 2G2RY.N.G and thus corresponds to GGRRRY.

20

**Table III**

R1	R2		R3		R4		R5	R6	R7
000	001		010		011		100	101	110
Step1(7)	N	R	G	GR	B	BR	BG	NOT USED:BGR	
Step2(6)	N	Y	2R	2RY	2G	2GY		NOT USED:2G2R, 2G2RY	
Step3(6)	N	2B	2Y	2Y2B	2X	2X2B			

30 Note: By convention, make the following replacements:  $2X \leftarrow 2G2R$ ,  $2X2B \leftarrow 2G2RY$

In contrast to the complex task of encoding reaction histories in a multi-step combinatorial synthesis, many applications require the distinction of only a limited set of chemistries. Simple color codes (SCC) can be constructed for this purpose. While not matching the encoding capacity of the corresponding binary color codes, these color codes are  
 5 entirely suitable in many instances in which the chemical distinctions of interest are created in a single reaction step, such as the coupling of a diagnostic probe to a bead. Examples of such limited chemical complexity include sensing applications as well as multi-agent monitoring and diagnostics.

As with binary color codes, the construction of simple color codes takes advantage of  
 10 distinguishable wavelengths, lifetimes and intensities of available fluorophores. A general version of the SCC based on a total of  $m$  fluorophores is constructed by using equal amounts of  $l$  fluorophores to encode each distinct chemical species of interest, where  $1 \leq l \leq m$ . In this code, the set of possible combinations of colors is equivalent to the number of possible configurations,  $S_r(l,m)$ , of a sample of size  $l$  drawn with replacement from a reservoir of  $m$ ,  
 15  $S_r(l,m) = (m+l-1)!/l!(m-l)!$ . Replacement allows for multiple instances of one color in each string.

For example, if 4 distinct fluorophores ( $m=4$ ) were available, and combinations of 3 ( $l=3$ ) were used - in equal relative abundances - for each distinct chemical species of interest, the generalized SCC would provide a total of 20 distinct configurations. These are listed in  
 20 table IV, denoting by R, G, B and Y the colors in a 4-color alphabet. Thus, the SCC ( $l=3$ ,  $m=4$ ) will uniquely encode the products generated in a single step of coupling up to 20 distinct antibodies to carrier beads; each of 20 reaction vessels would receive a mixture of three fluorophores in accordance with the set listed Table IV. The presence of several known fluorophores provides the basis to invoke coincidence methods to detect and monitor weak  
 25 signals and so to enhance assay sensitivity.

**Table IV**

	(R,R,R)	(G,G,G)	(B,B,B)	(Y,Y,Y)
30	(R,R,G)	(G,G,B)	(B,B,Y)	
	(R,R,B)	(G,G,Y)		



(R,R,Y)  
 (R,G,G)      (G,B,B)      (B,Y,Y)  
 (R,G,B)      (G,B,Y)  
 (R,G,Y)  
 5      (R,B,B)      (G,Y,Y)  
 (R,B,Y)  
 (R,Y,Y)

---

10      EXtended simple color codes (XSCC) can be constructed by varying relative abundances of fluorophores to create a set of distinguishable intensity levels for each of the fluorophore species in the alphabet. As with the XBCC, the XSCC permits control of  $m_i$  intensity levels for each of  $m_F$  fluorophore species in the alphabet.

15      Particularly easy to realize is the special case of SCC and XSCC where  $l=1$ ; only a single fluorophore marks each chemical species of interest.

### *Further Enhancements*

20      All color codes previously discussed herein can be further augmented by varying certain physico-chemical parameters of beads. For example, the number of encoded configurations may each be attached to a set of beads whose respective shapes, mean sizes, polarizabilities or other physico-chemical properties differ sufficiently so as to be distinguishable. By using  $S$  distinct sets of beads, the number of encoded configurations represented with XBCC( $m$ ) is increased to  $S \cdot 2^m$ .

25      BCC and XBCC encode chemical compound identity in terms of the relative abundances of fluorophores coupled to each bead. Accordingly, all permutations of a string of fluorophore tags are equivalent because they result in the same relative abundances. However, it has not escaped our notice that the implementation of the color code in which labeling leads to compound termination (see Fig. 3A) also retains a record of the order in which different color labels were added to each bead. Consequently, the analysis of molecular  
 30      weights of labeled compounds will reveal the order in which labeling occurred.

### *Chemical Realization of Extended Binary Color Code*

The realization of a chemical color code relies on a set ("alphabet") of chemically activated fluorophores with minimally overlapping absorption and emission spectra. We discuss here the case of the Extended Binary Color Code; other codes may be realized in analogous fashion. Although the implementation of a color code according to the present invention is illustrated herein by way of a specific family of fluorophores, the method is equally suitable for implementation with other fluorophores and chromophores whose distinctive spectral features serve to construct an alphabet of tags as described herein. An example of a suitable alphabet of six colors is provided by the CyDye(TM) family of indocyanine dyes, listed in Fig. 5.

The synthetic steps in this example are as follows (using standard Fmoc main-chain protection chemistry (Atherton & Sheppard, "Solid Phase Peptide Synthesis: A Practical Approach", IRL Press at Oxford University Press, Oxford, 1989, the contents are included herein by reference)).

**Table V**

- 
- 1) deprotect  $\alpha$ -amino group
  - 2) split resin population into a small number of aliquots
  - 3) for each resin aliquot, perform sub-stoichiometric coupling with coding CyDye activated ester;  
typical concentration:  $\approx 0.001$  to  $0.1$  mole of dye(s) per mole of  $\alpha$ -amino
  - 4) for each resin aliquot, perform coupling reaction with encoded amino acid
  - 5) pool resin aliquots
  - 6) repeat steps 1-5 for each randomized position in the amino acid sequence
- 

This procedure avoids fluorescence energy transfer between different dyes. First, labeling of any amino acid sequence as described herein will inactivate and so will terminate that sequence. Consequently, only a single dye is incorporated into any sequence and intra-sequence energy transfer is avoided. Second, low densities of dyes immobilized on the resin surface (see step 3 above) will ensure that lateral distances between labeled amino acid sequences substantially exceed the pertinent Förster radii for inter-strand fluorescent energy

transfer. This is a manifestation of the well known phenomenon of "pseudo-dilution" in solid phase synthesis.

The practicability of the procedure in Table V has been demonstrated by labeling standard combination synthesis bead resins (NovaSyn TG amino resin, NovaBiochem. "Combinatorial Chemistry" Catalog, San Diego, CA, 1997, the contents of which are included herein by reference). Specifically, we have constructed SCC( $l=1$ ,  $m=6$ ) as well as XSCC( $l=1$ ,  $m_F=1$ ,  $m_I=5$ ) with individual dyes and with multiple dyes of the CyDye series and have shown that colors are distinguishable by fluorescence microscopy at molar ratios as low as 0.0001. In addition, we have demonstrated that the dye coupling chemistry is compatible with protein synthesis as specified in Table V.

The method of the present invention may be used to realize color encoding of amino acid or peptide combinatorial libraries, examples of which are summarized in Table VI. A suitable reporter system is an anti- $\beta$ -endorphin monoclonal antibody (mAb) directed against an epitope in the form of an N-terminal amino acid sequence  $N_{tes}$ -YGGFL, where Y denotes tyrosine; binding of the primary anti- $\beta$ -endorphin mAb to its target is detected by a cascade-blue labeled secondary anti-mouse antibody (excitation at 396 nm, emission at 410 nm).

**Table VI**

20	<b>Binary Color Code (BCC)</b> bit 1: Cy2 bit 3: Cy5 bit 2: Cy3 bit 4: Cy7	XXGFL- $\beta$ Ala-BEAD X=Gly,Ala,Tyr,Phe	16=4x4 species created 16=2 <sup>4</sup> species created
25	<b>2-Level eXtended BCC</b> bit 1: Cy2 bit 5: Cy5 bit 2: 2*Cy2 bit 6: 2*Cy5 bit 3: Cy3 bit 7: Cy7 bit 4: 2*Cy3 bit 8: 2*Cy7	ZXXFL- $\beta$ Ala-BEAD Z=Gly,Ala,Glu,Lys, Phe,Tyr,D-Tyr X=Gly,Ala,Glu,Lys, Phe,Tyr	252=7*6*6 species created 256=2 <sup>8</sup> species encoded
30	<b>3-Level eXtended BCC</b> bit 1: Cy2 bit 7: Cy5 bit 2: 2*Cys2 bit 8: 2*Cy5 bit 3: 4*Cy2 bit 9: 4*Cy5 bit 4: Cy3 bit 10: Cy7 bit 5: 2*Cy3 bit 11: 2*Cy7 bit 6: 4*Cy3 bit 12: 4*Cy7	XXXXL- $\beta$ Ala-BEAD X=Gly,Ala,Ser,Asn, Glu,Lys,Phe,Tyr	4096=8 <sup>4</sup> species created 4096=2 <sup>12</sup> species encoded

Although the method of the present invention is illustrated by making reference to peptides and peptide precursors, the method is equally suitable with any other chemical precursors and compound classes that have been created via DCR combinatorial synthesis  
 5 (Calbiochem-NovaBiochem, "Solid Phase Organic Chemistry Handbook", San Diego, CA, 1997, the contents of which are included herein by reference).

Compounds prepared by the disclosed methods have potential use as therapeutic agents in the treatment of hypertension, inflammation, and analgesia. For example, enkephalin analogues selected by the disclosed methods may be useful as analgesics. Organic  
 10 compounds such as benzodiazepines useful as a muscle relaxant may also be selected by the disclosed methods.

#### *Diagnostics and Environmental Monitoring of Multiple Agents*

The method of the present invention enables a novel implementation of diagnostic  
 15 assays and tests that probe simultaneously for multiple reagents or pathogens. In contrast to the spatial encoding of diagnostic panels in all prior art, random assemblies of multiple bead types, distinguishable by their respective color codes, can be mixed and handled in parallel. For example, the implementation of bead-based immunodiagnostic assay formats can take advantage of color coding as described herein to display a multiplicity of specific bead-  
 20 anchored antibodies, each type assigned to a specific color code, to monitor for a multiplicity of agents in the ambient.

A preferred implementation of a multi-agent diagnostic assay uses random arrays of chemically encoded beads (Fig. 6). For example, the determination of blood type would require only five distinct bead types, a task that is readily addressed by the SCC ( $l=1$ ,  $m=5$ ).  
 25 This realization of diagnostic testing and environmental monitoring devices would facilitate miniaturization, integration of multiple tests and automated operation relying on spectral read-out.

#### *In-Situ Interrogation and Decoding of Color-Encoded Beads*

30 The optical arrangement in Fig. 7 provides for the integration of two essential capabilities: fluorescence microscopic imaging and multi-color fluorescence analysis of

individual beads. The latter serves to determine the relative abundances of several fluorophores present on the bead surface.

The use of a microscope objective of high numerical aperture (N.A. = 0.7)(702) serves to maximize collection efficiency as well as spatial resolution. The principal additional components of Fig. 7 are: a long-pass filter to reject stray excitation light (704), a dichroic beam splitter (706) to separate beams for image formation by the field lens (708) and spectral analysis via focusing of the light (by lens 710) on the slit aperture of a grating monochromator (712) or, alternatively (not shown), on the entrance pupil of an optical fiber that is coupled to a grating monochromator; multi-color spectra are recorded by a CCD array (714). Infinity-corrected optical components offer convenience of implementation.

While simple long pass filters have been employed in DNA sequencing applications to reject stray excitation light supplied at a single wavelength, interference filters can be designed to provide multiple narrow (10 nm) pass-bands at several emission wavelengths characteristic of the CyDye family of fluorophores discussed herein. Similar fabrication techniques may be applied to the dichroic mirror. These considerations are particularly relevant to an epi-fluorescence geometry, a special case of reflection microscopy.

Among the suitable instrumental realizations of recording spectral information from individual color-encoded beads or collections of color-encoded beads are flow cytometric analysis and multi-spectral imaging. The latter permits the collection of spectral information from individual or multiple beads in the field of view of a microscope or other imaging device, as considered in Fig. 7.

Methods suitable for multi-spectral imaging include: multiplexing of distinct wavelengths of incident and emitted light and illumination with a superposition of multiple wavelengths, followed by dispersive imaging by means of a grating or prism (see Fig. 7) or followed by interferometric analysis of emitted light.

The first method is readily implemented using matching optical pass-band filters; these are mounted in filterwheels and positioned in incident and emitted light paths of a microscope. The synchronized rotation of the two filterwheels will insert matching pairs of excitation and emission filters (a reflective geometry will also require a suitable dichroic mirror) into the light path, producing a repeating series of images at each of the distinct wavelengths selected

one of the filter/mirror combination. This principle is realized, for example, in the Fluorescence Imaging MicroSpectrophotometer developed by Kairos Scientific (Santa Clara, CA).

In the second method, distinct wavelengths for illumination are produced by a multi-pass band filter/mirror combination; a prism is inserted into the output path. This configuration facilitates the imultaneous spectral analysis of multiple beads located in a rectangular slice of the field of view of the microscope. Light emitted from beads within this slice is imaged onto the entrance slit of the prism and is decomposed into its spectral components. This principle is realized in the PARISS Imaging Spectrometer attachment developed by LightForm (Belle Meade, NJ). In the third method, light from the entire field of view is analyzed inteferometrically: a pellicle beamsplitter in the output path produces two (coherent) light beams which are reflected by a mirror and recombined. As the beamsplitter is rotated, a small difference in pathlength is introduced between the two light beams, resulting in interference fringes as the two beams are recombined. These fringes contain the entire spectral information contained in the light emiited from the field of view of a microscope (Garini et al, Bioimaging 4, 65-72 (1996)). That is, as the beamsplitter is rotated, a continuous spetrum is generated for every position within the field of view, resulting in a three-dimensional representation of the data. This principle is realized in the SpectraCube system developed and marketed by Applied Spectral Imaging (Carlsbad, CA). In contrast to the first method, the second and third methods generate a continuous spectrum, facilitating spectral classification of overlapping emission bands.

The arrangements in Fig. 8 provide for additional flexibility in rejecting stray light by spatially separating incident light and emitted light collection in transmission and rejection microscopy, as illustrated in Figs. 8A and 8B, respectively. In addition, the use of specially deigned multi-pass band interference filters in the output light path is again an option.

The demands on the sensitivity of the multi-color fluorescence detection system derive from the number of fluorophores of each color expected to be present on a selected bead. A bead of radius  $R$  and surface area  $A=4\pi R^2$  will accommodate up to  $N=A/a$  molecules of molecular area  $a$ , or  $N^*=xN$  fluorophores. With  $a=30A$  and  $0.01<x<0.1$ , a bead of 10  $\mu\text{m}$  diameter may carry  $10^7 \leq N^* \leq 10^8$  flurophores. For comparison, imaging of small circular domains of 10 $\mu\text{m}$  diameter within a monomolecular film composed of a phospholipid

containing 1 mole% of a fluorescent analog and confined to an air-water interface, is based on a comparable number of fluorophores and is readily accomplished using silicon-intensified target (SIT) camera technology. The refractive property of beads in aqueous solution will further enhance the light collection efficiency of the entire system.

5

### *In-situ Interrogation and Decoding of Color-Encoded Bead Arrays*

The present invention provides a methodology for color-encoding of beads and describes a method and apparatus for in-situ interrogation and decoding of color-encoded beads and collections of beads by multi-color fluorescence imaging and spectral analysis.

10 This method is compatible with all bead assay formats described to date, as discussed herein.

A preferred format providing a particularly efficient realization of bead assays on the basis of the methods and apparatus of the present invention involves planar beads arrays. This format facilitates highly parallel screening of enzyme activity, receptor-ligand binding, antibody-antigen recognition as well as DNA or RNA hybridization, etc. Thus, a close-packed array of 100 $\mu$ m diameter beads can contain of the order of  $10^4$  beads in an area of only 1 cm<sup>2</sup>, permitting the examination of up to  $10^4$  compounds/cm<sup>2</sup> in a single pass. The instantaneous determination of chemical identities enables the efficient implementation of re-iterative screening in which multiple copies of each bead type are examined to establish a statistically robust ranking of compounds producing positive assay scores. Furthermore, the implementation of the present invention in a planar bead array format lends itself to automation. Automated operation would entail the preparation of planar bead arrays, followed by fluorescence imaging of the array to locate beads that are to be subjected to spectral analysis and on-line decoding. The intrinsic detection sensitivity of fluorescence, demonstrated at the level of detecting single fluorophores, makes it possible to substantially reduce the size of synthesis beads. This in turn facilitates miniaturization and containment within an enclosed system, with its attendant benefits of reducing the requisite quantity of synthesized compound and the amount of reagents consumed in the course of screening.

15  
20  
25

One method of forming planar bead arrays is to rely on gravity-driven settling of beads from suspension to produce a (static) layer of beads or arrangement of bead clusters on a planar substrate. A second method employs dynamic planar bead arrays that are formed adjacent to planar surfaces and manipulated in-situ under external control, for example by

30

Light-controlled Electrokinetic Assembly of Particles near Surfaces (LEAPS). LEAPS is a technology that provides the capability to form dynamic planar bead arrays in aqueous solution on cue and to place and maintain them in a designated area of a planar electrode surface, as set forth in the copending PCT application filed April 24, 1997, entitled "Light  
 5 Controlled Electrokinetic Assembly of Particles Near Surfaces", based on U.S. Provisional Application Serial No. 60/016,642, filed April 25, 1996, which is incorporated by reference herein.

Dynamic planar bead arrays provide additional advantages in the realization of automated screening assays in a miniaturized, contained environment. Bead suspensions from  
 10 a synthesis pool will be loaded into a "sandwich" flow cell where planar bead arrays are formed adjacent to the planar walls of cell; screening assays will be performed in planar array format to identify lead compounds without the need of a time-consuming and error-prone step of physical separation; following completion of the scheduled assays, bead arrays will be disassembled and the bead suspension discharged to ready the flow cell for another cycle. In the  
 15 example, a redundancy of 10, i.e., the presence of 10 copies of beads of identical type and color code, would still facilitate screening of 1000 compounds at a time, but would considerably enhance the quality of any pharmacokinetic characterization. The benefits of miniaturization would be enhanced by the use of small synthesis beads. Chemically and physically well defined beads in the requisite size range (10 $\mu$ m diameter) are available from  
 20 many commercial sources. They are readily manipulated by LEAPS to form dynamic planar bead arrays of high density. This ensures that screening assays may be performed in a highly parallel format on a large number of samples, and this in turn provides the basis for highly re-iterative screening and for a robust pharmacokinetic characterization of potential lead compounds.

25 The present invention will be better understood from the Experimental Details which follow. However, one skilled in the art will readily appreciate that the specific methods and results discussed are merely illustrative of the invention as described in the claims which follow thereafter.

### Example 1

#### 30 1. Color-encoded PEG-polystyrene microspheres

##### a. Preparation of color-encoded PEG-polystyrene microspheres



- (1) Cy2 (ex = 489 nm, em = 506 nm)-color-encoded PEG-polystyrene microspheres:  
50 mg of NovaSyn TG amino microspheres (NovaBiochem; 130  $\mu$  diameter, 15  $\mu$ mol amine) were equilibrated in 10 ml DMF 30 min at 25°C. The supernatant was removed by filtration, and 100  $\mu$ l DMF, 1  $\mu$ l TEA and 15  $\mu$ l 1 mM Cy2-bisfunctional NHS-ester (Amersham; 15 nmol) were added in DMF. The reaction mixture was shaken 1 h at 25°C, 2  $\mu$ l (20  $\mu$ mole) n-butylamine was added, and the reaction mixture was shaken a further 30 min at 25°C. The supernatant was removed, and microspheres were washed twice with 5 ml DMF, rinsed twice with 5 ml chloroform and dried *in vacuo*.
- (2) Cy3 (ex = 550 nm, em = 570 nm)-color-encoded PEG-polystyrene microspheres:  
10 This preparation was identical to (1) except that, in parallel reactions, 15  $\mu$ l of 0.001, 0.01, 0.1, and 1 mM Cy3-monofunctional NHS-ester (Amersham; 0.15, 1.5, and 15 nmol) were used, and the n-butylamine step was omitted.
- (3) Cy3.5 (ex = 581 nm, em = 596 nm)-color-encoded PEG-polystyrene microspheres:  
This preparation was identical to (1) except that 15  $\mu$ l of 1 mM Cy3.5-monofunctional NHS-ester (Amersham; 15 nmol) was used, and the n-butylamine was step omitted.  
15
- (4) Cy5 (ex = 649 nm, em = 670 nm)-color-encoded PEG-polystyrene microspheres:  
This preparation was identical to (1) except that 15  $\mu$ l of 1mM Cy5-monofunctional NHS-ester (Amersham; 15 nmol) was used, and the n-butylamine step was omitted.
- (5) Cy5.5 (ex = 675 nm, em = 694 nm)-color-encoded PEG-polystyrene microspheres:  
20 This preparation was identical to (1) except that 15  $\mu$ l of 1 mM Cy5.5-monofunctional NHS-ester (Amersham; 15 nmol) was used, and the n-butylamine step was omitted.
- (6) Cy7 (ex = 743 nm, em = 767 nm)-color-encoded PEG-polystyrene microspheres:  
This preparation was identical to (1) except that 15  $\mu$ l of 1 mM Cy7-bisfunctional NHS-ester (Amersham; 15 nmol) was used.
- (7) Cy3/Cy5-color-encoded PEG-polystyrene microspheres:  
25 This preparation was identical to (1) except that both Cy3-monofunctional NHS-ester and Cy5-monofunctional NHS-ester were added (15  $\mu$ l of 1 mM stock each), and the n-butylamine step was omitted.
- (8) Cy2/Cy3/Cy5/Cy7-color-encoded PEG-polystyrene microspheres:

This preparation was identical to (1) except that Cy2-bisfunctional NHS-ester, Cy3-monofunctional NHS-ester, Cy5-monofunctional NHS-ester, and Cy7-bisfunctional NHS-ester were added (15  $\mu$ l of 1 mM stock each).

- b. Stability of Cy3-encoded PEG-polystyrene microspheres to solid-phase peptide synthesis conditions.

Cy3-encoded PEG-polystyrene microspheres were subjected to one cycle of solid-phase peptide synthesis. 50 mg microspheres and 5 mg Fmoc(Lys)Boc-OBT [prepared by reacting 94 mg Fmoc(Lys)Boc-OH (NovaBiochem; 0.2 mmol), 48 mg DCC (Aldrich; 0.22 mmol) and 27 mg HOBT (Aldrich; 0.2 mmol) in 2 ml DMF for 0.5 h at 25°C, centrifuging at 2000x g 5 min at 25°C, and using 100  $\mu$ l of the supernatant) in 100  $\mu$ l DMF were shaken 0.5 h at 25 °C. The microspheres were filtered, suspended in 100  $\mu$ l 20 % piperidine in DMF 15 min at 25°C, washed twice with 5 ml  $\text{CHCl}_3$ , and dried. The UV/VIS absorbance and fluorescence properties of the Cy3-encoded PEG-polystyrene microspheres were unchanged.

- c. Optical properties of color-encoded PEG-polystyrene microspheres

- Microspheres examined for their optical properties included:

Cy3 (ex = 550 nm, em = 570 nm)-color-encoded PEG-polystyrene microspheres of four different intensity levels, prepared as described in section a-(2) above by reacting beads with 0.001, 0.01, 0.1 and 1mM Cy3, are denoted b3-0001, b3-001, b3-01 and b3-1, respectively; as a group, all the Cy3-encoded PEG-polystyrene microspheres are denoted b3-x.

- Cy5 (ex = 649 nm, em = 670 nm)-color-encoded PEG-polystyrene microspheres, prepared as described in section a-(2) above by reacting beads with 1mM Cy5, are denoted b5-1;

Cy3/Cy5-color-encoded PEG-polystyrene microspheres, prepared as described in section a-(2) above by reacting beads with 1mM Cy3/Cy5, are denoted b35-1.

- An aliquot of dried microspheres was suspended in DMF and dispersed on a silicon wafer; DMF was evaporated by gentle heating. All subsequent observations were made in air.

#### (1) Fluorescence Imaging

Observations were made with a Zeiss UEM microscope equipped for epifluorescence; combinations of excitation filter/dichroic mirror/emission filter designed for Cy3 and Cy5 (Chroma Technologies, Brattleboro, VT) were used in conjunction with a 100W halogen

illuminator and objectives of 10X, 25X and 40X magnification. Optionally, images were recorded with a SIT camera (Cohu, San Diego, CA).

All microspheres displayed a bright circumferential “ring” of high intensity, corresponding to  $\leq 5\%$  of the particle diameter, suggesting that label was associated primarily with the surface, rather than the interior, of each particle. Even the dimmest particles, of type b3-0001, were readily observable using a 25X/0.45NA objective and the SIT camera. Microspheres of type b3-0001 appeared dimmer than did microspheres of type b3-001, although by less than the expected factor of 10. This phenomenon remains to be explored, but may indicate fluorescence quenching. Any given set of Cy3-encoded microspheres displayed particle-to-particle variations in color: some particles appeared orange, others yellow. of type b5-1 appeared bright red.

## (2) Fluorescence Spectra

To demonstrate the feasibility of in-situ interrogation of color-encoded microspheres, fluorescence spectra were recorded from individual color-encoded PEG-polystyrene microspheres by means of a PARISS<sup>TM</sup> imaging spectrophotometer (prototype supplied by LightForm, Belle Meade, NJ) with 50 $\mu$ m wide entrance slit, curved prism and room-temperature CCD array capable of on-chip integration. The instrument was mounted to the camera port of a Zeiss UEM microscope. In this configuration, multiple beads which are lined up along the long dimension of the projected slit can be imaged and spectrally analyzed. Only an approximate wavelength calibration was performed.

Spectra displaying fluorescence intensity as a function of wavelength were obtained separately for Cy3- and for Cy5-encoded microspheres and showed the following spectral characteristics:

b3-x: spectra were obtained for all types of particles; specific features included: for b3-0001: signal-to-noise (S/N)  $\approx 2$ , signal-to-background (S/B)  $\approx 1.5$ ; for b3-001: S/N  $\approx 4$ , S/B  $\approx 2$  (with a CCD integration time of approximately 10s); smoothing clearly revealed characteristic spectral features; for b3-1: S/N  $> 10$ ;  
b5-1: very clean spectra were recorded, all with a slight skew toward high wavelength;

b35-1: very clean spectra of either label were recorded, switching between appropriate filters to simulate filter wheel operation. At this concentration, spectra (taken with 10-times shorter integration time than that used for b3-01 and b3-001) displayed no discernible noise.

5

## 2. *Color-encoded macroporous polystyrene microspheres*

### a. Preparation of color-encoded macroporous polystyrene microspheres

50 mg Amino-Biolinker-PM1-1000 amino oligoethylene glycol-functionalized macroporous polystyrene microspheres (Solid Phase Sciences; 35  $\mu$  diameter, 7  $\mu$ mol amine) were equilibrated in 2 ml DMF 20 min at 25°C. The supernatant was removed by filtration, and 100  $\mu$ l DMF, 1  $\mu$ l TEA, and 70  $\mu$ l 1 mM Cy3-monofunctional NHS-ester (Amersham; 70 nmol) were added. After 1 hr at 25°C with shaking, the supernatant was removed by filtration, and the microspheres were washed twice with 5 ml DMF, washed twice with 5 ml  $\text{CHCl}_3$ , and dried *in vacuo*.

### 15 b. Optical properties of color-encoded macroporous polystyrene microspheres

Visual inspection using the configuration described under Example 1, revealed substantial bead-to-bead variations in fluorescence intensity.

## 3. *Color-encoded solid glass microspheres ("pelicular microspheres")*

### 20 a. Preparation of color-encoded pelicular microspheres

#### (1) Epoxide-functionalized pelicular microspheres:

4 g solid sodalime glass microspheres (Duke Scientific; 40 $\pm$ 3  $\mu$  diameter; 4.8 x 10<sup>7</sup> microspheres), 7 ml xylene, 2.34 ml 3-glycidoxypropyltrimethoxysilane (Aldrich; 1 mmol) and 0.117 ml diisopropylethylamine (Aldrich; 0.7 mmol) were shaken 18 h at 80°C. Upon cooling to room temperature, microspheres were filtered, washed with 40 ml methanol, washed with 40 ml diethyl ether, and dried *in vacuo*.

#### (2) MMT-NH-PEG-functionalized pelicular microspheres:

Microspheres from (1) were suspended in a solution of 200 mg mono-MMT-1,13-trioxotridecadiamine [0.4 mmol; prepared by mixing 7 g MMT-Cl (Aldrich; 23 mmol) and 11.3 ml 4,7,10-trioxa-1,13-tridecanediamine (Aldrich; 51 mmol) in 150 ml 1:1:1 methylene chloride:pyridine:acetonitrile for 18 h at 25°C, then isolating the required adduct by

30

chromatography on silica gel) in 6 ml xylene. Approximately 10 mg sodium hydride (Aldrich; 0.4 mmol) was added, and the suspension shaken 18 h at 40°C under a drying tube. Microspheres then were filtered and successively washed with 20 ml methanol, 10 ml water, 20 ml methanol, and 20 ml chloroform, and dried *in vacuo*.

- 5 Dried microspheres were capped by reaction with 5% acetic anhydride, 5% 2,6-lutidine, 8% N-methylimidazole in 10 ml tetrahydrofuran 1 h at 25°C with shaking, successively washed in 2x5 ml methanol, 2x5 ml chloroform, and 2x5 ml diethyl ether, and dried *in vacuo*.

(3) H<sub>2</sub>N-PEG-functionalized pelicular microspheres:

Microspheres from (2) were treated with 1 ml 3% TFA in CH<sub>2</sub>Cl<sub>2</sub> 0.5 h at 25°C with shaking.

- 10 Based on quantitation of released monomethoxy trityl cation ( $\epsilon_{478} = 3.47 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ ) the loading densities of H<sub>2</sub>N-PEG were as follows:

15 15 fmol H<sub>2</sub>N-PEG per microsphere

1.1 x 10<sup>10</sup> molecules H<sub>2</sub>N-PEG per microsphere

0.022 molecule H<sub>2</sub>N-PEG per Å<sup>2</sup>

- 15 Assuming  $\approx 0.04$  available silanol groups per Å<sup>2</sup> of soda-lime glass, the grafting efficiency was  $\approx 50\%$ .

(4) Color-encoded PEG-functionalized pelicular microspheres:

- To 20 mg of H<sub>2</sub>N-PEG-functionalized pelicular microspheres (4.2 nmol amine), were added 97 µl DMF, 2 µl TEA, and 0.8 µl 1 mM Cy3-monofunctional NHS-ester (Amersham; 0.8  
20 nmol), and the resulting suspension was shaken for 18 h at 25°C. Microspheres then were filtered and washed successively with 5 ml DMF, 5 ml methanol, 5 ml chloroform, and 5 ml diethyl ether, and dried *in vacuo*.

Based on quantitation of consumed Cy3-monofunctional NHS-ester ( $\epsilon_{552} = 1.5 \times 10^5 \text{ M}^{-1} \text{ cm}^{-1}$ ) the loading of Cy3 densities were as follows:

- 25 1 fmol Cy3 per microsphere

6x10<sup>8</sup> molecules Cy3 per microsphere

0.001 molecule Cy3 per Å<sup>2</sup>

0.07 molecule Cy3 per molecule available H<sub>2</sub>N-PEG

b. Optical properties of Cy3-encoded PEG-functionalized pelicular microspheres:

- 30 Visual inspection using the configuration described under Example 1, revealed uniformly fluorescent microspheres.

**What is Claimed is:**

1. A method of identifying a compound having a selected property of interest in  
5 a library of compounds, each of said compounds being bound to its respective  
solid support, and being produced by a unique reaction series composed of  $N$   
reaction steps, wherein each compound is prepared from a component, and  $N$   
is an integer from at least 1 to about 100, which comprises:
- 10 a) dividing a population of solid supports having at least one type of a first  
functional group at the surface of said solid support selected from the  
group consisting of  $\text{CO}_2\text{H}$ ,  $\text{OH}$ ,  $\text{SH}$ ,  $\text{NH}_2$ ,  $\text{NHR}$ ,  $\text{CH}_2\text{Cl}$ ,  $\text{CH}_2\text{Br}$  and  $\text{CHN}_2$ ,  
wherein  $R$  is a linear  $\text{C}_1$ - $\text{C}_9$  alkyl group, into  $M$  batches, wherein  $M$  is an  
integer from at least 2 to about 25;
- 15 b) coupling the  $M$  batches of solid support in a set of at least one reaction  
respectively with  $M$  different components so as to form a bond with the  
solid support via said first functional group, said components being  
independently protected or unprotected;
- 20 c) adding to each batch, either prior to coupling step b), concurrently  
therewith, or subsequently to step b), from about 0.001 to about 0.5 molar  
equivalent of a spectrally distinguishable fluorophore tag associated  
uniquely with each component, said tag being identified by its  
characteristic excitation wavelength(s), emission wavelength(s), excited  
state lifetime and emission intensity, said tag being activated so as to be  
25 capable of forming either a direct bond to the surface of the solid support,  
either via the first or a second functional group which is protected or  
unprotected and is the same as or different from the first functional group  
bonded to the component, or an indirect bond via a  $\text{C}_1$ - $\text{C}_9$  linear or  
branched alkyl linker moiety which is either interrupted or uninterrupted  
by at least one oxygen or nitrogen atom or a carbonyl,  $(\text{C}=\text{O})\text{NH}$  or

NH(C=O) moiety, wherein when said second functional group is protected, said functional group is deprotected prior to forming said direct or indirect bond, said linker being bonded to the second functional group at the surface of the solid support; and either

- 5           d) recombining all *M* batches, said recombining step being either prior to or subsequent to step e) and steps e) - g); or
- e) performing an assay capable of indicating that any compound in the library either while bound to or cleaved from its solid support has the property of interest;
- 10          f) collecting spectral fluorescence data for each respective solid support so as to determine respective relative abundances of the fluorophore tags bound thereto; and
- g) analyzing the collected spectral fluorescence data by comparing the respective relative abundances of the fluorophore tags determined in step
- 15          f) so as to determine the unique reaction series for the compound, thereby identifying the compound having the property of interest.
2. The method of claim 1 wherein the components are independently selected from the group consisting of an amino acid, a hydroxyacid, an oligoamino acid, an
- 20          oligopeptide, a saccharide, an oligosaccharide, a diamine, a dicarboxylic acid, an amine-substituted sulfhydryl, a sulfhydryl-substituted carboxylic acid, an alicyclic, an aliphatic, a heteroaliphatic, an aromatic and a heterocyclic moiety.
3. The method of claim 2 wherein the saccharide is a suitably protected D- or L-
- 25          glucose, fructose, inositol, mannose, ribose, deoxyribose or fucose.
4. The method of claim 2 wherein the oligopeptide is an enkephalin, a vasopressin, an oxytocin, an atrial natrietic factor, a bombesin, a calcitonin, a parathyroid hormone, a neuropeptide Y or an endorphin, or a fragment thereof

comprising at least 20% of the components thereof, or an isosteric analogue thereof wherein independently  $\text{NH}(\text{C}=\text{O})$  is replaced by  $\text{NH}(\text{C}=\text{O})\text{NH}$ ,  $\text{NH}(\text{C}=\text{O})\text{O}$ ,  $\text{CH}_2(\text{C}=\text{O})$  or  $\text{CH}_2\text{O}$ ;  $\text{NH}_2$  is replaced by  $\text{OH}$ ,  $\text{SH}$ ,  $\text{NO}_2$  or  $\text{CH}_3$ ;  $\text{CH}_3\text{S}$  is replaced by  $\text{CH}_3(\text{S}=\text{O})$  or  $\text{CH}_3\text{CH}_2$ ; indole is replaced by naphthyl or indene; hydroxyphenyl is replaced to tolyl, mercaptophenyl or nitrophenyl; and/or hydrogen in an aromatic ring is replaced by chlorine, bromine, iodine or fluorine;  $\text{C}_1\text{-C}_4$  alkyl is replaced by partially or fully fluorinated  $\text{C}_1\text{-C}_4$  alkyl.

- 5  
10  
15  
20  
25
5. The method of claim 2 wherein the oligopeptide is an ACE inhibitor, an HIV protease inhibitor, a cytolytic oligopeptide or an antibacterial oligopeptide.
6. The method of claim 2 wherein the aromatic is para-disubstituted benzene, biphenyl, naphthalene or anthracene, either substituted or unsubstituted by linear or branched chain lower alkyl, alkoxy, halogen, hydroxy, cyano or nitro.
7. The method of claim 2 wherein the heterocyclic moiety is 2,6-disubstituted pyridine, thiophene, 3-7-disubstituted N-protected indole or 2,4-disubstituted imidazole, either substituted or unsubstituted by linear or branched chain lower alkyl, alkyl, halogen, hydroxy, cyano or nitro.
8. The method of claim 1 wherein the solid support is a microsphere, a bead, a resin or a particle, and is composed of a material selected from the group consisting of polystyrene, polyethylene, cellulose, polyacrylate, polyacrylamide, or preferably a silica to glass bead.
9. The method of claim 1 wherein the solid support is chemically modified by covalent attachment of either a substituted or unsubstituted oligo- or polyethyleneglycol, which either terminated or unterminated by an amine substituted by either hydroxymethyl, chloromethyl, aminomethyl or



mercaptomethyl, wherein the functional group at the surface of the solid support is hydroxy, chlorine,  $\text{NH}_2$  or  $\text{SH}$ , respectively.

5 10. The method of claim 1 wherein the assay is performed while the compound is cleaved from its solid support under conditions whereby the compound remains adsorbed to the solid support.



10 11. The method of claim 1 wherein the property of interest is binding affinity of a compound to a receptor, the assay is performed by determining a physical response to binding by

- 15 a) first admixing with the library of compounds a solution of a labelled receptor so as to result in labelled receptor bound to at least one compound bound to a solid support;
- b) removing the solution from the solid support; and either
- c) washing the solid support so as substantially to remove non-bound labelled receptor, and step (d), or
- d) measuring the physical response due to bound labelled receptor so as to determine the binding affinity.





20 12. The method of claim 11 wherein the receptor is labelled by a fluorescent dye, a colored dye, radioisotope or an enzyme.

13. The method of claim 11 wherein the physical response is fluorescence emission, optical absorption or radioactivity.



25 14. The method of claim 1 wherein the components have a structure independently selected from the group consisting of:



5



10



5 wherein R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub> and R<sub>6</sub> are independently methyl, ethyl, linear or branched chain C<sub>3</sub>-C<sub>9</sub>, phenyl, benzyl, benzoyl, cyano, nitro, halo, formyl, acetyl and linear or branched chain C<sub>3</sub>-C<sub>9</sub> acyl; wherein a, b, c, d and e are independently 0, 1, 2 or 3; wherein X, Y and Z are independently NH, O, S, S(=O), CO, (CO)O, O(CO), NH(C=O) or (C=O)NH; and wherein W is independently N, O or S.

- 10 15. The method of claim 1 wherein at least one component is an amino acid, bearing a protected or unprotected group which is capable of participating in a further reaction or coupling step and is nitrogen, said protecting group being selected from the group consisting of N-*α*-fluorenylmethyloxycarbonyl, t-butyloxycarbonyl, t-amylloxycarbonyl, (trialkylsilyl) ethyloxycarbonyl, t-butyl and benzyl.
- 15 16. The method of claim 1 wherein the fluorophore tag represents a bit of a binary code, and comprises zero, one or more than one fluorescent dye, multiple fluorescent dyes, said dye(s) being spectrally distinguishable by excitation wavelength, emission wavelength, excited-stated lifetime or emission intensity.
- 20 17. The method of claim 16 wherein emission intensity is distinguished by adjusting the ration of the relative quantities of each fluorophore.
18. The method of claim 17 wherein the ratio is 1:1, 2:1; 3:1 or 4:1.
- 25 19. The method of claim 1 wherein the fluorophore tags are dyes selected from the group consisting of compounds with the chemical names:
  - 3-( $\epsilon$ -carboxypentyl)-3'-ethyl-oxacarbocyanine-6,6'-disulfonic acid
  - 1-( $\epsilon$ -carboxypentyl)-1'-ethyl-3,3,3',3'-tetramethylindocarbocyanine-5,5'-disulfonic acid

1-( $\epsilon$ -carboxypentyl)-1'-ethyl-3,3,3',3'-tetramethyl-3H-

benz(e)indocarbocyanine-5,5',7,7'-tetrasulfonic acid

1-( $\epsilon$ -carboxypentyl)-1'-ethyl-3,3,3',3'-tetramethylindocarbocyanine-5,5'-

disulfonic acid

5

1-( $\epsilon$ -carboxypentyl)-1'-ethyl-3,3,3',3'-tetramethyl-3H-

benz(e)indodicarbocyanine-5,5',7,7'-tetrasulfonic acid

1-( $\epsilon$ -carboxypentyl)-1'-ethyl-3,3,3',3'-tetramethylindotricarbocyanine-5,5'-

disulfonic acid

and are activated as active esters selected from the group consisting of succinimidyl, sulfosuccinimidyl, p-nitrophenol, pentafluorophenol, HOBt and N-hydroxypiperidyl.

10

20. The method of claim 1 wherein the fluorophore tags are dyes selected from the group consisting of compounds with the chemical names:

15

6-((4,4-difluoro-5,7-dimethyl-4-bora-3a,4a-diaza-s-indacene-

3-propionyl)amino) hexanoic acid

6-((4,4-difluoro-5-phenyl-4-bora-3a,4a-diaza-s-indacene-3-propionyl)

amino) hexanoic acid,

20

6-((4,4-difluoro-1,3-dimethyl-5-(4-methoxyphenyl)-4-bora-3a,

4a-diaza-s-indacene-2-propionyl) amino)hexanoic acid,

6-(((4-(4,4-difluoro-5-(2-thienyl)-4-bora-3a,4a-diaza-s-indacene-3-yl)

phenoxy) acetyl) amino)hexanoic acid,

6-(((4-(4,4-difluoro-5-(2-thienyl)-4-bora-3a,4a-diaza-s-indacene-3-yl)

styryloxy)acetyl) aminohexanoic acid, and

25


6-(((4-(4,4-difluoro-5-(2-pyrrolyl)-4-bora-3a,4a-diaza-s-indacene-3-yl)

styryloxy) acetyl)aminohexanoic acid,


and are activated as active esters selected from the group consisting of succinimidyl, sulfosuccinimidyl, p-nitrophenol, pentafluorophenol, HOBt and N-hydroxypiperidyl.

21. The method of claim 1 wherein the fluorophore tags are dyes selected from the group consisting of compounds with the chemical structures:


5



10





15



20

25





and are activated as active esters selected from the group consisting of succinimidyl, sulfosuccinimidyl, p-nitrophenol, pentafluorophenol, HOBt and N-hydroxypiperidyl.

22. The method of claim 1 wherein the assay is performed by cleaving compounds from the solid support while permitting diffusion through solution and binding to receptors, said receptors being arranged in proximity to each solid support.

23. The method of claim 1 wherein the fluorescence data are collected from multiple solid supports using multi-spectral imaging methods.

24. The method of claim 1 wherein one of the fluorophore tags uniquely associated with a preselected component or reaction comprises a ligand and a substance capable of binding specifically to the ligand, said ligand being labelled with a fluorophore and attached in a post-assay reaction, said substance being present on the solid support and attached prior to, concurrently with, or subsequent to the coupling of the component, whereby the labelled ligand when bound to the substance indicates the presence of the preselected component.

25. The method of claim 1 wherein the solid support is a polymeric bead, and spectral fluorescence data is collected by
- a) forming either a static planar array or a dynamic planar array of beads; and
  - b) obtaining a fluorescence image for each bead.

5

26. The method of claim 25 wherein the planar array of beads is formed adjacent to the planar walls of a sandwich flow cell and controlled by light-controlled electrokinetic means.

10

27. The method of claim 25 wherein the planar array of beads is formed by using an apparatus capable of dynamically assembling and disassembling an array of beads at an interface between an electrode and an electrolyte solution, said apparatus comprising:

15

- i) an electrode, an electrolyte solution and an interface therebetween
- ii) a plurality of beads located in said electrolyte solutions;
- iii) said electrode being patterned to include at least one area of modified electrochemical properties;
- iv) an illumination source which illuminates said electrode with a predetermined light pattern;
- 20 v) an electric field generator which generates an electric field at said interface to cause the assembly of an array of beads in accordance with the predetermined light pattern and the electrochemical properties of said electrode; and
- 25 vi) an electric field removal unit which removes said electric field to cause the disassembly of said array of beads.

28. The method of claim 25 wherein spectral fluorescence data are collected for the bead array by initially forming a spatially encoded array of beads

suspended at an interface between an electrode and an electrolyte solution, comprising the following steps:

- 5
- ii) providing multiple types of particles, each type being stored in accordance with chemically or physically distinguishable particle characteristics in one of a plurality of reservoirs, each reservoir containing a plurality of like-type particles suspended in said electrolyte solution;
- 10
- iii) providing said reservoirs in the form of an  $m \times n$  grid arrangement;
- iv) patterning said electrode to define  $m \times n$  compartments corresponding to said  $m \times n$  grid of reservoirs;
- 15
- v) depositing  $m \times n$  droplets from said  $m \times n$  reservoirs onto said corresponding  $m \times n$  compartments, each said droplet originating from one of said reservoirs and remaining confined to one of said  $m \times n$  compartments and each said droplet containing at least one particle;
- 20
- vi) positioning a top electrode above said droplets so as to simultaneously contact each said droplet;
- vii) generating an electric field between said top electrode and said  $m \times n$  droplets;
- 25
- viii) using said electric field to form a bead array in each of said  $M \times N$  compartments, each said bead array remaining spatially confined to one of said  $m \times n$  droplets;
- ix) illuminating said  $m \times n$  compartments on said patterned electrode with a predetermined light pattern to maintain the position of said bead arrays in accordance with said predetermined light and the pattern of  $m \times n$  compartments; and
- x) positioning said top electrode closer to said electrode thereby fusing said  $m \times n$  droplets into a continuous liquid phase, while



maintaining each of said  $m \times n$  bead arrays in one of the corresponding  $m \times n$  compartments.

- 5           29. The method of claim 28, wherein said compartments are hydrophilic and the remainder of said electrode surface is hydrophobic.
30. The method of claim 1 wherein  $N$  is an integer from at least 2.
- 10          31. The method of claim 1 wherein  $N$  is an integer from at least 4 to about 12.
32. The method of claim 1 wherein  $M$  is an integer from at least 4 to about 10
33. The method of claim 1 wherein from about 0.01 to about 0.05 molar equivalent of a spectrally distinguishable fluorophore tag is added in step c).
- 15          34. A compound having a selected property of interest as identified in accord with claim 1.
35. A chemical library prepared in accord with claim 1.
- 20          36. An apparatus for identifying a compound having a selected property of interest in a library of compounds, each of said compounds being bound to its respective solid support, and being produced by a unique reaction series composed of  $N$  reaction steps, wherein each said compound is prepared from a component, and  $N$  is an integer from at least 1 to about 100, said solid support being at least one particle array, said apparatus comprising:
- 25           a) an electrode and an electrolyte solution having an interface therebetween,
- b) an electric field generator which generates an electric field at an interface between an electrode and an electrolyte solution;

- 5 c) said electrode being patterned to modify the electrochemical properties of said electrode;
- d) an illuminating source which illuminates said interface with a predetermined light pattern to control the movement of said particles in accordance with said predetermined light pattern and the electrochemical properties of said electrode;
- 10 e) means for preparing said chemical library, which comprises:
- i) means for dividing a population of solid supports having at least one type of a first functional group at the surface of said solid support selected from the group consisting of  $\text{CO}_2\text{H}$ ,  $\text{OH}$ ,  $\text{SH}$ ,  $\text{NH}_2$ ,  $\text{NHR}$ ,  $\text{CH}_2\text{Cl}$ ,  $\text{CH}_2\text{Br}$  and  $\text{CHN}_2$ , wherein R is a linear  $\text{C}_1$ - $\text{C}_9$  alkyl group, into  $M$  batches, wherein  $M$  is an integer from at least 2 to about 25;
- 15 ii) means for coupling the  $M$  batches of solid support in a set of at least one reaction respectively with  $M$  different components so as to form a bond with the solid support via said first functional group, said components being independently protected or unprotected;
- 20 iii) means for adding to each batch either prior to coupling step ii), concurrently therewith, or subsequently to step ii), from about 0.001 to about 0.5 molar equivalent of a spectrally distinguishable fluorophore tag associated uniquely with each component, said tag being identified by its characteristic excitation wavelength(s), emission wavelength(s), excited state lifetime and emission intensity, said tag being activated so as to be capable of forming
- 25 either a direct bond to the surface of the solid support, either via the first or a second functional group which is protected or unprotected and is the same as or different from said first functional group, a direct bond to the component which if

5                   protected is priorly deprotected, or an indirect bond via a C<sub>1</sub>-C<sub>9</sub>,  
linear or branched alkyl linker moiety which is either interrupted  
or uninterrupted by at least one oxygen or nitrogen atom or a  
carbonyl, (C=O)NH or NH(C=O) moiety, wherein when said  
second functional group is protected, said second functional group  
is deprotected prior to forming said direct or indirect bond, said  
linker being bonded to said second functional group at the surface  
of the solid support; and either

- 10                   iv)       means for recombining all *M* batches, said recombining step either  
being prior to or subsequent to step v), and steps v)-vii); or;
- v)       means for performing an assay capable of indicating that any  
compound in the library either while bound to or cleaved from its  
solid support has the property of interest;
- 15                   vi)       means for collecting spectral fluorescence data for each respective  
solid support so as to determine respective relative abundances of  
the fluorophore tags bound thereto;
- vii)       means for analyzing the collected spectral fluorescence data by  
comparing the respective relative abundances of the fluorophore  
tags determined in step vi) so as to determine the unique reaction  
series for the compound, thereby identifying the compound having  
20                   the property of interest.

25                   37. A method of identifying a compound having a selected property of interest in  
a library of compounds, each of said compounds being bound to its respective  
solid support, and being produced by a unique reaction series composed of *N*  
coupling or reaction steps, wherein each compound is prepared from  
components which are independently the same or different, and *N* is an integer  
from at least 1 to about 100, which comprises:

- 5 a) dividing a population of solid supports having at least one type of a first functional group at the surface of said solid support surface selected from the group consisting of  $\text{CO}_2\text{H}$ ,  $\text{OH}$ ,  $\text{SH}$ ,  $\text{NH}_2$ ,  $\text{NHR}$ ,  $\text{CH}_2\text{Cl}$ ,  $\text{CH}_2\text{Br}$  and  $\text{CHN}_2$ , wherein R is a linear  $\text{C}_1\text{-C}_9$  alkyl group, into  $M$  batches, wherein  $M$  is an integer from at least 2 to about 50;
- 10 b) coupling the  $M$  batches of solid support in a set of at least one reaction respectively with  $M$  different initial components so as to form a bond with the solid support via said first functional group, said components being protected or unprotected at a group which is capable of participating in a further coupling step and orthogonally protected at non-participating group(s);
- 15 c) adding to each batch either prior to coupling step b), concurrently therewith, or subsequently to step b), from about 0.001 to about 0.5 molar equivalent of a spectrally distinguishable fluorophore tag associated uniquely with each initial component or a reaction of step b), said tag being identified by its characteristic excitation wavelength(s), emission wavelength(s), excited state lifetime and emission intensity, said tag being activated so as to be capable of forming either a direct bond to the surface of the solid support, either via the first or a second functional group which is protected or unprotected and is the same as or different from said first functional group, a direct bond to the initial component which if protected is priorly deprotected, or an indirect bond via a  $\text{C}_1\text{-C}_9$  linear or branched alkyl linker moiety which is interrupted or uninterrupted by either at least one oxygen or nitrogen atom or a carbonyl,  $(\text{C}=\text{O})\text{NH}$  or  $\text{NH}(\text{C}=\text{O})$  moiety, said linker being bonded to said first functional group at the surface of the solid support, wherein when said second functional group is protected, said second functional group is deprotected prior to forming said direct or indirect bond; and either
- 20
- 25

d) recombining all  $M$  batches and cleaving any protecting group present at a group which is to participate in a further coupling step, said recombining step being either prior to or subsequent to step e), and steps e)-h); or

e) iteratively  $N - 1$  times

- 5 (1) dividing a population of solid supports into  $M(N)$  batches, wherein  $M(N)$  depends on  $N$  and is an integer from at least 2 to about 25;
- 10 (2) coupling the  $M(N)$  batches of solid support respectively with  $M(N)$  different components, wherein  $M(N)$  is the number of batches during the  $N$ th step, said components being protected or not protected at a group which is capable of participating in a further coupling step and orthogonally protected at a nonparticipating group(s);
- 15 (3) adding to each batch either prior to coupling step (2), concurrently therewith, or subsequently to step (2), from about 0.001 to about 0.5 molar equivalent of a spectrally distinguishable fluorophore tag associated uniquely with each component in the  $N$ th coupling step (2), said tag being identified by its characteristic excitation wavelength(s), emission wavelength(s), excited state lifetime and emission intensity, said tag being activated so as to form either a
- 20 direct bond to the surface of the solid support, either via a functional group which is protected or not protected and is the same as or different from the functional group bonded to the component, a direct bond to the  $(N - 1)$ th component, or an indirect bond via a  $C_1$ - $C_9$  linear or branched alkyl linker moiety
- 25 which is optionally interrupted by at least one oxygen or nitrogen atom or a carbonyl,  $(C=O)NH$  or  $NH(C=O)$  moiety, said linker being bonded to the functional group at the surface of the solid support, wherein when said functional group is protected, said

function group is deprotected prior to forming said direct or indirect bond; and

(4) recombining all  $M(N)$  batches and cleaving any protecting group present at a group which is to participate in a further coupling

5 step;

so as to form a compound having  $N$  components;

f) performing an assay capable of indicating that any compound in the library either while bound to or cleaved from its solid support has the property of interest;

10 g) collecting spectral fluorescence data for each respective solid support so as to determine respective relative abundances of the fluorophore tags bound thereto; and

h) analyzing the collected spectral fluorescence data by comparing the respective relative abundances of the fluorophore tags determined in step

15 g) so as to determine the  $N$  components coupled in the unique reaction series for the compound, thereby identifying the compound having the property of interest.

20 38. The method of claim 37 wherein the components are independently selected from the group consisting of an amino acid, a hydroxyacid, an oligoamino acid, an oligopeptide, a saccharide, an oligosaccharide, a diamine, a dicarboxylic acid, an amine-substituted sulfhydryl, a sulfhydryl-substituted carboxylic acid, an alicyclic, an aliphatic, a heteroaliphatic, an aromatic and a heterocyclic moiety.

25 39. The method of claim 38 wherein the saccharide is a suitably protected D- or L-glucose, fructose, inositol, mannose, ribose, deoxyribose or fucose.

40. The method of claim 38 wherein the oligopeptide is an enkephalin, a vasopressin, an oxytocin, an atrial natrietic factor, a bombesin, a calcitonin, a parathyroid hormone, a neuropeptide Y or an endorphin, or a fragment thereof comprising at least 20% of the components thereof, or an isosteric analogue thereof wherein independently  $\text{NH}(\text{C}=\text{O})$  is replaced by  $\text{NH}(\text{C}=\text{O})\text{NH}$ ,  $\text{NH}(\text{C}=\text{O})\text{O}$ ,  $\text{CH}_2(\text{C}=\text{O})$  or  $\text{CH}_2\text{O}$ ;  $\text{NH}_2$ , is replaced by  $\text{OH}$ ,  $\text{SH}$ ,  $\text{NO}_2$ ,  $\text{CH}_3$ ;  $\text{CH}_3$ ,  $\text{S}$  is replaced by  $\text{CH}_3$ ,  $(\text{S}=\text{O})$  or  $\text{CH}_3$ ,  $\text{CH}_2$ ; indole is replaced by naphthyl or indene; hydroxyphenyl is replaced by tolyl, mercaptophenyl or nitrophenyl; and/or hydrogen in an aromatic ring is replaced by chlorine, bromine, iodine or fluorine;  $\text{C}_1$ - $\text{C}_4$  alkyl is replaced by partially or fully fluorinated  $\text{C}_1$ -,  $\text{C}_4$ -, alkyl.
41. The method of claim 38 wherein the oligopeptide is an ACE inhibitor, an HIV protease inhibitor, a cytolytic oligopeptide or an antibacterial oligopeptide.
42. The method of claim 38 wherein the aromatic is para-di substituted benzene, biphenyl, naphthalene or anthracene, either substituted or unsubstituted by linear or branched chain lower alkyl, alkoxy, halogen, hydroxy, cyano or nitro.
43. The method of claim 38 wherein the heterocyclic moiety is 2,6-disubstituted pyridine, thiophene, 3,7-disubstituted N-protected indole or 2,4-disubstituted imidazole, either substituted or unsubstituted by linear or branched chain lower alkyl, alkoxy, halogen, hydroxy, cyano or nitro.
44. The method of claim 37 wherein the solid support is a microsphere, a bead, a resin or a particle, and is composed of a material selected from the group consisting of polystyrene, polyethylene, cellulose, polyacrylate, polyacrylamide, or preferably a silica or glass bead.



- 5 45. The method of claim 37 wherein the solid support is chemically modified by covalent attachment of a substituted or unsubstituted oligo- or polyethyleneglycol, which is either terminated or unterminated by an amine substituted by either hydroxymethyl, chloromethyl, aminomethyl or mercaptomethyl, wherein the functional group at the surface of the solid support is hydroxy, chlorine,  $\text{NH}_2$  or  $\text{SH}$ , respectively.
- 10 46. The method of claim 37 wherein the assay is performed while the compound is attached to its solid support.
47. The method of claim 37 wherein the assay is performed while the compound is cleaved from its solid support under conditions whereby the compound remains adsorbed to the solid support.
- 15 48. The method of claim 37 wherein when the property of interest is binding affinity of a compound to a receptor, the assay is performed by determining a physical response to binding by
- 20 a) first admixing with the library of compounds a solution of a labelled receptor so as to result in labelled receptor bound to at least one compound bound to a solid support;
- b) removing the solution from the solid support; and either
- c) washing the solid support so as substantially to remove non-bound labelled receptor, and step (d); or
- 25 d) measuring the physical response due to bound labelled receptor so as to determine the binding affinity.
49. The method of claim 48 wherein receptor is labelled by a fluorescent dye, a colored dye, radioisotope or an enzyme.







50. The method of claim 48 wherein the physical response is fluorescence emission, optical absorption or radioactivity.



51. The method of claim 37 wherein the components have a structure independently selected from the group consisting of:

5



10





- 5 wherein  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ , and  $R_6$  are independently methyl, ethyl, linear or branched chain  $C_3$ - $C_9$  alkyl, phenyl, benzyl, benzoyl, cyano, nitro, halo, formyl, acetyl and linear or branched chain  $C_3$ - $C_9$  acyl; wherein  $a$ ,  $b$ ,  $c$ ,  $d$  and  $e$  are independently 0, 1, 2 or 3; wherein  $X$ ,  $Y$  and  $Z$  are independently NH, O, S, S(=O), CO, (CO)O, O(CO), NH(C=O) or (C=O) NH; and wherein  $W$  is
- 10 independently N, O or S.

52. The method of claim 37 wherein at least one component is an amino acid, and the protected or unprotected group which is to participate in a further coupling step is nitrogen, said protecting group being selected from the group consisting of  $N$ - $\alpha$ -fluorenylmethyloxycarbonyl,  $t$ -butyloxycarbonyl,  $t$ -amyloxycarbonyl, (trialkylsilyl) ethyloxycarbonyl,  $t$ -butyl and benzyl;
- 15
53. The method of claim 37 wherein the fluorophore tag represents a bit of binary code, and comprises zero, one or more than one fluorescence dye, multiple fluorescent dyes, said dye(s) being spectrally distinguishable by excitation wavelength, emission wavelength, excited-state lifetime or emission intensity.
- 20

5

54. The method of claim 37 wherein the assay is performed by cleaving compounds from the solid support while permitting diffusion through solution and binding to receptors, said receptors being arranged in proximity to each solid support.

10

55. The method of claim 37 wherein the fluorescence data are collected from multiple solid supports using multi-spectral imaging methods.

56. The method of claim 53 wherein emission intensity is distinguished by adjusting the ratio of the relative quantities of each fluorophore.

57. The method of claim 56 wherein the ratio is 1:1, 2:1, 3:1 or 4:1.

15

58. The method of claim 37 wherein the fluorophore tags are dyes selected from the group consisting of compounds with the chemical names:

3-( $\epsilon$ -carboxypentyl)-3'-ethyl-oxacarbocyanine-6,6'-disulfonic acid

1-( $\epsilon$ -carboxypentyl)-1'-ethyl-3,3,3',3'-tetramethylindocarbocyanine-5,5'-disulfonic acid

20

1-( $\epsilon$ -carboxypentyl)-1'-ethyl-3,3,3',3'-tetramethyl-3H-

benz(e)indocarbocyanine-5,5',7,7'-tetrasulfonic acid

1-( $\epsilon$ -carboxypentyl)-1'-ethyl-3,3,3',3'-tetramethylindocarbocyanine-5,5'-disulfonic acid

25

1-( $\epsilon$ -carboxypentyl)-1'-ethyl-3,3,3',3'-tetramethyl-3H-

benz(e)indodicarbocyanine-5,5',7,7'-tetrasulfonic acid

1-( $\epsilon$ -carboxypentyl)-1'-ethyl-3,3,3',3'-tetramethylindotricarbocyanine-5,5'-disulfonic acid

and are activated as active esters selected from the group consisting of succinimidyl, sulfosuccinimidyl, p-nitrophenol, pentafluorophenol, HOBt and N-hydroxypiperidyl

59. The method of claim 37 wherein the fluorophore tags are dyes selected from the group consisting of compounds with the chemical names:

6-((4,4-difluoro-5,7-dimethyl-4-bora-3a,4a-diaza-s-indacene-3-propionyl)amino) hexanoic acid

6-((4,4-difluoro-5-phenyl-4-bora-3a,4a-diaza-s-indacene-3-propionyl) amino) hexanoic acid,

6-((4,4-difluoro-1,3-dimethyl-5-(4-methoxyphenyl)-4-bora-3a,4a-diaza-s-indacene-2-propionyl) amino)hexanoic acid,


6-(((4-(4,4-difluoro-5-(2-thienyl)-4-bora-3a,4a-diaza-s-indacene-3-yl) phenoxy) acetyl) amino)hexanoic acid,

6-(((4,4-difluoro-5-(2-thienyl)-4-bora-3a,4a-diaza-s-indacene-3-yl) styryloxy)acetyl) aminohexanoic acid, and

6-(((4,4-difluoro-5-(2-pyrrolyl)-4-bora-3a,4a-diaza-s-indacene-3-yl) styryloxy) acetyl)aminohexanoic acid,


and are activated as active esters selected from the group consisting of succinimidyl, sulfosuccinimidyl, p-nitrophenol, pentafluorophenol, HOBt and N-hydroxypiperidyl.

60. The method of claim 37 wherein the fluorophore tags are dyes selected from the group consisting of compounds with the chemical structures:




50


5



10




15



20

25



and are activated as active esters selected from the group consisting of succinimidyl, sulfosuccinimidyl, p-nitrophenol, pentafluorophenol, HOBt and N-hydroxypiperidyl.

- 5 61. The method of claim 37 wherein one of the fluorophore tags uniquely associated with a preselected component or reaction comprises a ligand and a substance capable of binding specifically to the ligand, said ligand being labelled with a fluorophore and attached in a post-assay reaction, said substance being present on the solid support and attached prior to, concurrently with, or subsequent to the coupling of the component, whereby the labelled ligand when bound to the substance indicates the presence of the preselected component.
- 10
- 15 62. The method of claim 37 wherein the solid support is a bead, and spectral fluorescence data are collected by
- a) forming either a static planar array or a dynamic planar array of beads; and
  - b) obtaining a fluorescence image for at least one bead.
- 20 63. The method of claim 62 wherein the planar array of beads is formed adjacent to the planar walls of a sandwich flow cell and controlled by light-controlled electrokinetic means.
- 25 64. The method of claim 62 wherein the dynamic planar array of beads is formed by using an apparatus capable of dynamically assembling and disassembling an array of beads at an interface between an electrode and an electrolyte solution, said apparatus comprising:
- i) an electrode, an electrolyte solution and an interface there between;
  - ii) a plurality of beads located in said electrolyte solution;

- iii) said electrode being patterned to include at least one area of modified electrochemical properties;
- iv) an illumination source which illuminates said electrode with a predetermined light pattern;
- 5 v) an electric field generator which generates an electric field at said interface to cause the assembly of an array of beads in accordance with the predetermined light pattern and the electrochemical properties of said electrode; and
- 10 vi) an electric field removal unit which removes said electric field to cause the disassembly of said array of beads.
65. The method of claim 62 wherein spectral fluorescence data are collected for the bead array by initially forming a spatially encoded array of beads suspended at an interface between an electrode and an electrolyte solution,
- 15 comprising the following steps:
- i) providing an electrode and an electrolyte solution;
- ii) providing multiple types of particles, each type being stored in accordance with chemically or physically distinguishable particle characteristics in one of a plurality of reservoirs, each reservoir containing a plurality of like-type particles suspended in said
- 20 electrolyte solution;
- iii) providing said reservoirs in the form of an  $m \times n$  grid arrangement;
- iv) patterning said electrode to define  $m \times n$  compartments corresponding to said  $m \times n$  grid of reservoirs;
- 25 v) depositing  $m \times n$  droplets from said  $m \times n$  reservoirs onto said corresponding  $m \times n$  compartments, each said droplet originating from one of said reservoirs and remaining confined to one of said  $m \times n$  compartments and each said droplet containing at least one particle;

- 5 vi) positioning a top electrode above said droplets so as to simultaneously contact each said droplet;
- vii) generating an electric field between said top electrode and said  $m \times n$  droplets;
- 10 viii) using said electric field to form a bead array in each of said  $m \times n$  compartments, each said bead array remaining spatially confined to one of said  $m \times n$  droplets;
- ix) illuminating said  $m \times n$  compartments on said patterned electrode with a predetermined light pattern to maintain the position of said bead arrays in accordance with said predetermined light pattern and the pattern of  $m \times n$  compartments; and
- 15 x) positioning said top electrode closer to said electrode thereby fusing said  $m \times n$  droplets into a continuous liquid phase, while maintaining each of said  $m \times n$  bead arrays in one of the corresponding  $m \times n$  compartments.
66. The method of claim 65 wherein said compartments are hydrophilic and the remainder of said electrode surface is hydrophobic.
- 20 67. The method of claim 37 wherein  $N$  is an integer from at least 3 to about 12.
68. The method of claim 37 wherein  $M$  and  $M(N)$  are independently an integer from at least 4 to about 12.
- 25 69. The method of claim 37 wherein from about 0.01 to about 0.05 molar equivalent of a spectrally distinguishable fluorophore tag is added in step c).
70. A compound having a selected property of interest as identified in accord with claim 37.



71. A chemical library prepared in accord with claim 37.

72. An apparatus for identifying a compound having a selected property of interest in a library of compounds, each of said compounds being bound to its respective solid support, and being produced by a unique reaction series composed of  $N$  coupling and reaction steps, wherein each said compound is prepared from a set of components which are independently the same or different, and  $N$  is an integer from at least 1 to about 100, said solid support being at least one particle array, said apparatus comprising:

- a) an electrode and an electrolyte solution having an interface therebetween;
- b) an electric field generator which generates an electric field at an interface between an electrode and an electrolyte solution;
- c) said electrode being patterned to modify the electrochemical properties of said electrode;
- d) an illuminating source which illuminates said interface with a predetermined light pattern to control the movement of said particles in accordance with said predetermined light pattern and the electrochemical properties of said electrode;
- e) means for preparing said chemical library, which comprises:
  - i) means for dividing a population of solid supports having at least one type of a first functional group at the surface of said solid support selected from the group consisting of  $\text{CO}_2\text{H}$ ,  $\text{OH}$ ,  $\text{SH}$ ,  $\text{NH}_2$ ,  $\text{NHR}$ ,  $\text{CH}_2\text{Cl}$ ,  $\text{CH}_2\text{Br}$  and  $\text{CHN}_2$ , wherein  $R$  is a linear  $\text{C}_1\text{-C}_9$  alkyl group, into  $M$  batches, wherein  $M$  is an integer from at least 2 to about 50;
  - ii) means for coupling the  $M$  batches of solid support in a set of at least one reaction respectively with  $M$  different initial components so as to form a bond with the solid support via said first functional group, said components being protected or unprotected at a group which is

to participate in a further coupling step and orthogonally protected at non-participating group(s);

- iii) means for adding to each batch either prior to coupling step ii), concurrently therewith, or subsequently to step ii), from about 0.001 to about 0.5 molar equivalent of a spectrally distinguishable fluorophore tag associated uniquely with each initial component, said tag being identified by its characteristic excitation wavelength(s), emission wavelength(s), excited state lifetime and emission intensity, said tag being activated so as to be capable of forming either a direct bond to the surface of the solid support, either via the first or a second functional group which is protected or unprotected and is the same as or different from said first functional group bonded to the component, or an indirect bond via a  $C_1$ - $C_9$ , linear or branched alkyl linker moiety which is either interrupted or uninterrupted by either at least one oxygen or nitrogen atom or a carbonyl,  $(C=O)NH$  or  $NH(C=O)$  moiety, said linker being bonded to said second functional group at the surface of the solid support, wherein when said second functional group is protected, said second functional group is deprotected prior to forming said direct or indirect bond; and either
- iv) means for recombining all  $M$  batches and cleaving any protecting group present at a group which is to participate in a further coupling step, and steps v)-viii); or
- v) means for iteratively  $N - 1$  times
- (1) dividing a population of solid supports into  $M(N)$  batches, wherein  $M(N)$  depends on  $N$  and is an integer from at least 2 to about 25;
  - (2) coupling the  $M(N)$  batches of solid supports respectively with  $M(N)$  different components, wherein  $M(N)$  is the number of

batches during the  $N$ th step, said components being protected or unprotected at a group which is capable of participating in a further coupling step and orthogonally protected at a non-participating group(s);

- 5 (3) adding to each batch either prior to coupling step (2), concurrently therewith, or subsequently to step (2), from about 0.001 to about 0.1 molar equivalent of a different spectrally distinguishable fluorophore tag associated uniquely with each component during the  $N$ th coupling step (2), said tag being uniquely identified by its excitation wavelength, emission wavelength, excited-state lifetime or emission intensity, whereby said tag is activated so as to be capable of forming either a direct bond to the solid support, either via an  $N$ th functional group which is protected or unprotected and is the same as or different from the first functional group, or an indirect bond thereto via a  $C_1$ - $C_9$  linear or branched alkyl linker moiety which is either interrupted or uninterrupted by either at least one oxygen or nitrogen atom or a carbonyl or  $NH(C=O)$  moiety, or a direct bond to the  $(N-1)$ th component which if protected is priorly deprotected, said tag or linker being bound via the group which is to participate in a further coupling step, wherein when said  $N$ th functional group is protected, said  $N$ th functional group is deprotected prior to forming said direct or indirect bond; and
- 10
- 15
- 20
- 25 (4) recombining all  $M(N)$  batches and cleaving the protecting group present if present at a group which is to participate in a further coupling step;

so as to form a compound having  $N$  components;

- 5
- 10
- vi) means for performing an assay capable of indicating that any compound in the library either while bound to or cleaved from its solid support has the property of interest;
  - vii) means for collecting spectral fluorescence data for each respective solid support so as to determine respective relative abundances of the fluorophore tags bound thereto;
  - viii) means for analyzing the collected spectral fluorescence data by comparing the respective relative abundances of the fluorophore tags determined in step vii) so as to determine the  $N$  components coupled in the unique reaction series for the compound, thereby identifying the compound having the selected property of interest.

# Divide-Couple-Recombine(DCR) Combinatorial Synthesis





FIGURE 1

[illegible]

**SHEET 2 OF 9**

### Strategies for Tag-Coupling





FIGURE 3

# COLOR-CODING OF PCR COMBINATORIAL SYNTHESIS

## Step 1: Divide beads into 4 groups




## Couple with first set of amino acids and color labels




## Recombine beads

## Step 2: Divide beads into 4 groups



## Couple with second set of amino acids and color labels




## Recombine beads

FIGURE 4



## Emission Spectra of the CyDye Fluorescent Dyes



Fluorophore	Color Of Fluorescence	Absorption Maximum (nm)	Fluorescence Maximum (nm)	Extinction Coefficient (M <sup>-1</sup> cm <sup>-1</sup> )	Quantum Yield for protein conjugates	Formula Weight (daltons)	
						Bisfnc.	Monofnc.
Cy2	Green	489 nm	506 nm	~ 150,000	>0.12	896.95	713.78
Cy3	Orange	550 nm	570 nm	150,000	>0.15	949.11	765.95
Cy3.5	Scarlet	581 nm	596 nm	150,000	>0.15	1285.54	1102.37
Cy5	Far-Red	649 nm	670 nm	250,000	>0.28	975.15	791.99
Cy5.5	Near IR	675 nm	694 nm	250,000	>0.28	1311.58	1128.41
Cy7	Near IR	743 nm	767 nm	~ 250,000	~ 0.28	1001.19	818.02
FluorX	Green	494 nm	520 nm	68,000	0.3		586.6

**FIGURE 5**

# **Multi-Agent Diagnostic Assay In Bead-Array Format**



Transparent Cover




Beads in Buffer



Planar Substrate

Not to Scale


**Side View**



**RandomBead  
Array (5 Colors)**

**Top View**

**FIGURE 6**



**FIGURE 7**

# Multicolor Fluorescence Imaging and Analysis




FIGURE 8




FIGURE 9